

Pamphlet to accompany:

NCGS Open File Report 2026-03

Compiled bedrock geologic map of the Southern Pines 30' x 60' Quadrangle (Southern Pines 100K), North Carolina. (version 9/1/2026)

New data collection and compilation work by: Philip J. Bradley, Garrett J. Thompson, Qingxin Zhang, Douglas B. Hill, Ezra T. Becker, and Katherine E. Pelt

Cartographic representation by Garrett J. Thompson, Philip J. Bradley, and Michael A. Medina

New geochronology data reduction and interpretation by Adam C. Curry

2026

Reference:

Bradley, P.J., Thompson, G.J., Zhang, Q., Becker, E.T., Hill, D.B., Pelt, K.E., and Curry, A.C., 2026, Compiled Bedrock Geologic Map of the Southern Pines 30' x 60' Quadrangle (Southern Pines 100K), North Carolina, North Carolina Geological Survey Open-file Report 2026-03, scale 1:100,000, in color, with accompanying pamphlet.

BACKGROUND

This compiled bedrock geologic map, partially supported by the U.S. Geological Survey (USGS) National Cooperative Geologic Mapping Program under STATEMAP, completes year 2 of a 2-year project to compile the entire bedrock portion of the Southern Pines 30'x 60' Quadrangle (Southern Pines 100K) in support of the USGS US Geoframework Initiative's (USGI) vision for a nation-wide, seamless geologic map.

The goal of this compilation effort is to produce a new 1:100,000-scale digital geologic map of the study area using the USGS Geologic Mapping Schema (GeMS). When available, geologic data at scales more detailed than 1:100,000-scale were used (i.e. 1:24,000-scale data). The FY24 (May 2024 to June 2025) and FY25 (July 2025 to August 2026) phases of the compilation effort compiled legacy data at scales ranging from 125K- to 24K-scale in the northern and southern parts of the Southern Pines 100K, respectively. NCGS staff conducted targeted foot and vehicle traverses to validate contacts from legacy sources and to collect new field data. Close attention was paid to rectifying edge-match issues between legacy data and recent 24K-scale mapping to allow accurate transitions of map units from areas with detailed data to areas of legacy data collected at smaller scales (e.g. 48K). More detailed and closer-spaced traverses targeted areas of structural or stratigraphic complexity. Less detailed and wider spaced traverses were used in areas of less complex structure or rock types (e.g. areas underlain by homogenous plutons received less attention). LiDAR data displayed as hillshade was used to help identify lineaments along with possible faults and dikes that may be groundwater sources or preferred pathways for groundwater contaminants.

PAST WORKERS

Specific data sources are listed in the DataSources table in the geodatabase. Major data sources are provided in the index of workers figure on the map layout.

GEOLOGIC SETTING

The Southern Pines 100K includes multiple major geologic elements (see figure on map). From west to east, they are: the Carolina terrane, Deep River Triassic basin (Sanford- and Wadesboro-sub-basins), the Carolina terrane east of the Wadesboro sub-basin, the Lilesville, Millstone Lake, and newly identified Pinebluff plutons, the Ellerbe Triassic basin, the easternmost Carolina terrane, the newly identified Upper Little River terrane, and large areas of Coastal Plain sedimentary deposits. Bedrock of the Crabtree and Spring Hope terranes are interpreted to be present in the southeastern portion of the map area and buried by Coastal Plain sedimentary deposits. Older alluvium deposits are present along major drainages in addition to modern floodplain deposits. This map is a compiled bedrock map; surficial sediments are not displayed.

Interpretation of Sub-Coastal Plain Geology

Large portions of the Southern Pines 100K are overlain by Atlantic Coastal Plain sediments – see inset figure on geologic map. For areas in which the bedrock is covered by Coastal Plain sediments, data from the NC Department of Transportation Geotechnical Engineering Unit, boring data (NCDOT, 2026) and Lawrence and Hoffman (1993) - Geology of basement rocks beneath the North Carolina Coastal Plain - were utilized. The sub-Coastal Plain geology of Lawrence and Hoffman (1993) was based on widely spaced drilled data and geophysical data that included aeromagnetic maps (Zietz et al., 1984) and Bouguer gravity maps (Black, 1986; Daniels and Leo, 1985). Additionally, the locations of some diabase dikes and terrane boundaries were modified from Lawrence and Hoffman (1993) after review of the preliminary magnetic survey map from the Piedmont Bridge South Magnetic and Radiometric Survey (USGS Piedmont Bridge, 2025). Additional edits to the sub-Coastal Plain geology in the map area will likely occur after the official release and review of the new Piedmont Bridge data.

CAROLINA TERRANE

The Carolina terrane underlies the northwest portion of the map area. The Carolina terrane is composed of Neoproterozoic to Cambrian metamorphosed volcanic, volcano-sedimentary, sedimentary and intrusive rocks (Hibbard et al., 2002; and Hibbard et al., 2006). The Carolina terrane is separated into three lithotectonic units: 1) the Hyco arc, 2) the Aaron Formation of the redefined Virgilina sequence (Hibbard et al., 2013), and 3) the Albemarle arc (Hibbard et al., 2013).

COMPILED GEOCHRONOLOGY DATES FOR THE CAROLINA TERRANE IN THE SOUTHERN PINES 100K

As part of the compilation efforts within the Southern Pines 100K, crystallization age and detrital zircon age data was compiled. Additionally, through STATEMAP support, new samples were collected and analyzed. Geochronologic data is summarized in Table 1 with references. Sample locations are identified and labeled on the geologic map. Geochronology plots and tables of NCGS collected samples are provided in Appendix A.

Table 1: Summary of available age dates in the Southern Pines 100K. Refer to main map for sample locations. Geochronology plots and tables of NCGS collected samples are provided in Appendix A.

Label	Sample ID	Rock Type	Unit	Date	Source	Comments
A	SB-9	rhyolitic tuff	Albemarle Arc-Uwharrie Volcanic Complex	554 ± 15 Ma weighted average of six $^{206}\text{Pb}/^{238}\text{U}$ ages for zircon grains by SHRIMP methods	U-Pb zircon crystallization age (Ingle et al., 2003)	Location resampled by NCGS – See sample B.
B	NCGS Station SP100K-550027	rhyodacite	Albemarle Arc-Uwharrie Volcanic Complex	537.6 ± 2.8 Ma (MSWD = 1.8)	U-Pb zircon dates by AZ LaserChron with data interpretation by A. Curry (2025), preliminary data	Abner Mountain area. Resample of Ingle et al. (2003) SB-9.
C	NCGS Station SP100K-550264	rhyodacite	Albemarle Arc-Morrow Mountain Rhyodacite	553.2 ± 2.5 Ma (MSWD = 1.6)	U-Pb zircon dates by AZ LaserChron with data interpretation by A. Curry (2025), preliminary data	Troy landfill area
D	NCGS Station SP100K-550946	tuffaceous pebbly sandstone	Albemarle Arc	MDA - ca. 524 Ma	Detrital zircon dates (AZ LaserChron). Data interpretation by A. Curry (2025), preliminary data	Seagrove formation of the Robbins area - volcaniclastic sedimentary rock
E	NCGS Station SP100K-550363	conglomeratic sandstone	Albemarle Arc	MDA - ca. 543 Ma	Detrital zircon dates (AZ LaserChron). Data interpretation by A. Curry (2025), preliminary data	Seagrove formation of the Biscoe area - volcaniclastic sedimentary rock

F	Lemon Springs Quarry sample	Metagranite	Easternmost Carolina terrane	Rb-Sr date of 429+/-10 Ma	Kish and Campbell, 1986	Minimum age. Rock is likely older.
G	NCGS Station SP100K-551063	rhyodacite	Albemarle Arc-Morrow Mountain Rhyodacite	550.4 ± 4.2 Ma (MSWD = 0.44)	U-Pb zircon dates by AZ LaserChron with data interpretation by A. Curry (2025), preliminary data	Rhyodacite dome in Robbins area
H	NCGS-SP100K 551983	Laminated siltstone	Albemarle arc	MDA - ca. 540 Ma	Detrital zircon dates (AZ LaserChron). Data interpretation by A. Curry (2026), preliminary data	Ellerbe area volcanic and volcanoclastic rocks
I	NCGS-SP100K 551641	metamorphosed siltstone	Albemarle arc	MDA - ca. 562 Ma	Detrital zircon dates (AZ LaserChron). Data interpretation by A. Curry (2026), preliminary data	Ellerbe area volcanic and volcanoclastic rocks
J	NCGS-SP100K 551786	metamorphosed sandy siltstone	Albemarle arc	MDA - ca. 543 Ma	Detrital zircon dates (AZ LaserChron). Data interpretation by A. Curry (2026), preliminary data	Ellerbe area volcanic and volcanoclastic rocks

HYCO ARC AND THE HYCO FORMATION

In the map area and adjacent Chapel Hill 100K, the Hyco Arc consists of the Hyco Formation which includes ca. 633 to 612 Ma (Wortman et al., 2000; Bowman, 2010; Bradley and Miller, 2011 and Bowman et al., 2013) metamorphosed layered volcanoclastic rocks and plutonic rocks. Dating by Barefoot (2015), indicates that magmatism in the lower portions of the Hyco Formation may be as old as ca. 650 Ma. Available age dates (Wortman et al., 2000; Bradley and Miller, 2011) indicate the Hyco Formation may tentatively be divided into lower (ca. 630 Ma) and upper (ca. 615 Ma) portions with an apparent intervening hiatus of magmatism.

Map units of metavolcanic and metavolcanoclastic rocks include various lithologies that, when grouped together, are interpreted to indicate general environments of deposition. The dacitic lavas and tuffs unit is interpreted to represent dacitic domes and proximal pyroclastics. The andesitic to basaltic lavas (with tuffs or conglomerates) units are interpreted to represent eruption of intermediate to mafic lava flows

and associated pyroclastic and/or epiclastic deposits. The epiclastic/pyroclastic units are interpreted to represent deposition from the erosion of dormant and active volcanic highlands. Deposition of the primary volcanics ranged from subaerial to subaqueous with concomitant deposition of various types and volumes of volcanic-derived sediments. Some of the metavolcanic units within the map area display lithologic relationships similar to dated units present in northern Orange and Durham Counties. Due to these similarities, the metavolcanic and metavolcaniclastic units have been tentatively separated into upper and lower portions of the Hyco Formation; geochronologic data in the map area is needed to confirm this interpretation.

Generally, Hyco Formation volcanic and sedimentary rocks are quartz crystal *poor* or quartz is absent in hand sample. In contrast (discussed in a later section), the Seagrove formation sedimentary rocks and Uwharrie Volcanic Complex (Uwharrie Formation) volcano-sediments and volcanic rocks range from quartz crystal-rich to having quartz locally present.

Within the Southern Pines 100K, Hyco Formation related units outcrop in three locations: 1) The main outcrop area of the Hyco Formation is present in the north-central portion of the map in the High Falls area (north of Robbins). This area is the southwest extension of the bulk of the Hyco Formation that forms continuous map units north and northeast to the VA-NC State line. 2) Northeast of Troy is an outlier of Hyco Formation rocks that is disconnected from the main outcrop pattern. This area marks the northeast end of the Troy anticline. Complexly deformed rocks that are locally hydrothermally altered are present. The area is home to the Cottonstone Mountain hydrothermal deposit, the Star mine, and the Carter mine. The Hyco units are likely part of a dome structure with a high-angle reverse fault marking part of the map unit's eastern boundary. 3) West of Jugtown is another outlier of Hyco Formation rocks that was likely exhumed in an anticline that is disrupted by a thrust fault.

ALBEMARLE ARC UNITS

Age relationships between rocks mapped as the Uwharrie Formation (Conley and Bain, 1965; Stromquist and Sundelius, 1969) and the Albemarle Group (Milton, 1984) are unclear. Traditionally, the Uwharrie Formation has been viewed as being the older substrate to the Albemarle Group (e.g. Butler and Secor, 1991); this interpretation is supported by recent models of intra-arc rifting (Moye, 2023). However, radiometric ages from rocks mapped as Uwharrie Formation suggest that rocks of the Uwharrie Formation and Albemarle Group overlap in age, forming a time transgressive sequence (Pollock et al., 2010 and Hibbard et al., 2013). Before the mapping and new geochronology of this study, the stratigraphic relationships of the rocks yielding younger-than-expected ages compared to the bulk of the Uwharrie Formation were not confidently established. The goals of this new mapping and compilation work were to compile existing maps, rectify edge-match issues, and collect rock samples for modern geochronologic analysis to better define the Uwharrie Formation and its relationship to adjacent units.

Stratigraphically, the main outcome of this new study is the redefinition of the Uwharrie Formation to the Uwharrie Volcanic Complex - a package of lithodemic units that includes metamorphosed: extrusive and shallow intrusive volcanic rocks, volcanoclastic pyroclastic rocks, volcanoclastic sedimentary rocks and sedimentary rocks. Additionally, as part of this study, the Seagrove formation is established and the

nomenclature of the Morrow Mountain Rhyodacite Suite is adopted (discussed below in more detail) (Figures 1 and 2). Refinement of the stratigraphy is ongoing.

MORROW MOUNTAIN RHYODACITE SUITE

The Morrow Mountain Rhyodacite Suite is a suite of intrusions with a range of ages and textures that are likely magmatically linked (Boorman et al., 2013 and Moye, 2023b). Isolated bodies of rhyodacite surrounded by volcano-sedimentary rocks were grouped as part of the suite. Texturally similar bodies are present in areas mapped as part of the rhyodacitic lavas and tuffs unit of the Uwharrie Volcanic Complex (discussed below).

A summary of existing and new age dates within the Southern Pines 100K map are provided in Table 1. For a summary of existing and new age dates in the Chapel Hill 100K, see the Open-file report of the Chapel Hill 100K (Bradley et al., 2025a).

Ingle et al. (2003) reported TIMS $^{207}\text{Pb}/^{206}\text{Pb}$ ages of 539 ± 5 Ma for the Morrow Mountain rhyodacite in the Charlotte 100K. Bradley et al. (2025b), reported LA-ICP-MS $^{206}\text{Pb}/^{238}\text{U}$ weighted mean ages for two samples in the Chapel Hill 100K: A sample from the Seagrove area yielded an age of 536.4 ± 6.1 Ma (MSWD = 4.5) and a sample from Caraway Mountain yielded an age of 542.7 ± 5.3 Ma (MSWD = 0.5).

New and unpublished LA-ICP-MS $^{206}\text{Pb}/^{238}\text{U}$ weighted mean ages completed in 2025 from samples from the suite includes: 548.8 ± 2.8 Ma (MSWD = 1.4) for a sample collected from Daves Mountain in Asheboro in the Chapel Hill 100K (Bradley et al., 2025a); 550.4 ± 4.2 Ma (MSWD = 0.44) for a sample collected from the Robbins area in the Southern Pines 100K (geochronology data location G); and 553.2 ± 2.5 Ma (MSWD = 1.6) for a sample collected near the landfill in the Troy area in the Southern Pines 100K (geochronology data location C). Two samples were analyzed from the eastern portion of the Charlotte 100K in the vicinity of Dennis Mountain in the Uwharrie National Forest. LA-ICP-MS $^{206}\text{Pb}/^{238}\text{U}$ weighted means ages from the samples yielded: 539.4 ± 1.8 Ma (MSWD = 1.6) (sample CLT100k_BoormanE7_CGS6) and 546.0 ± 3.1 Ma (MSWD = 1.6) (sample CLT100k_BoormanE9_CGS7).

The previously existing and new age data indicate that volcanism of the Morrow Mountain Rhyodacite Suite spanned from ca. 553 Ma to ca. 536 Ma and was contemporaneous with the Uwharrie Volcanic Complex (discussed below) and deposition in the Albemarle Group.

REDEFINITION OF THE UWHARRIE FORMATION TO UWHARRIE VOLCANIC COMPLEX

Seiders (1981) separated the Uwharrie Formation into multiple units that included metamorphosed felsic lavas, dikes, and volcanic necks of varying textures along with associated coarse-grained volcanoclastic, pyroclastic, and sedimentary rocks. The volcanoclastic rocks include metamorphosed tuffaceous sandstones, conglomeratic sandstones, and siltstones with lesser amounts of fine- to coarse tuff. Fiamme-like clasts are common in the conglomerates, sandstones and tuffs. Quartz and feldspar crystal fragments are common in the sedimentary components, tuffs and lavas. Subordinate amounts of

intermediate to mafic volcanics and volcano-sedimentary rocks and fine grained clastics (siltstones and mudstones) are also present in the Uwharrie Formation units of Seiders (1981).

Previous geochronological data indicated that the Uwharrie Formation is ca. 550 Ma (Ingles et al., 2003). Pollock et al. (2010), reported a youngest detrital zircon age of ca. 545 Ma from a sample reportedly from the oldest portions of the Formation. Hibbard et al. (2013) speculated that the Uwharrie Formation may be time transgressive or previous mapping was in error and that additional geochronologic and mapping work is needed to properly determine the extent and nature of the Uwharrie Formation. Recent geochronologic data from rocks assigned to the Uwharrie Formation indicate that problems exist with the definition of the Formation. Detrital zircon and crystallization ages from this study indicate that units previously interpreted as intrusive into surrounding volcanoclastic units are disconformable (e.g. - Daves Mountain in the Asheboro area in the Chapel Hill 100K is older than the surrounding clastic unit). Additionally, a sample of rhyodacite from the “core” of the Uwharrie Formation in the Chapel Hill 100K yields a crystallization age of ca. 537 Ma (Bradley, et al., 2025a) which is inconsistent with the area being the oldest portions of the Formation. According to the North American Stratigraphic Code, a Formation should have a defined stratigraphic position. Geochronologic data indicates that rocks mapped as part of the Uwharrie Formation are packages of rocks that represent volcanic centers and related volcanoclastic debris that vary in age. As such, the Uwharrie Formation may be better defined as a package of lithodemic units. While additional investigation into the stratigraphic nomenclature continues, for this map, the Formation nomenclature is dropped and Uwharrie Volcanic Complex is used to emphasize the volcanic origin and restricted regional distribution of the package of rocks.

Uwharrie Volcanic Complex (Uwharrie Formation) legacy and new geochronology data

A summary of existing and new age dates within the Southern Pines 100K map are provided in Table 1. For a summary of existing and new age dates in the Chapel Hill 100K, see the Open-file report of the Chapel Hill 100K (Bradley et al., 2025a).

Ingles et al. (2003) reported the weighted average of six $^{206}\text{Pb}/^{238}\text{U}$ ages for zircon grains by SHRIMP methods of 554 ± 15 Ma for a sample from the Uwharrie Formation (SB-9) in unit Zurlt in the Southern Pines 100K (geochronology data location A). This location was resampled as part of this study and yielded LA-ICP-MS $^{206}\text{Pb}/^{238}\text{U}$ weighted means ages of 537.6 ± 2.8 Ma (MSWD = 1.8) (geochronology data location B). Goliber and Coleman (2023) reported a CA-ID-TIMS weighted mean age of 548.7 ± 1.1 Ma from a felsic volcanic rock in the unit in the Chapel Hill 100K. A new and unpublished LA-ICP-MS $^{206}\text{Pb}/^{238}\text{U}$ weighted mean age yielded 537.4 ± 5.7 Ma (MSWD = 0.86) for a sample collected from Luther Place trail area in the Uwharrie Volcanic Complex in the Chapel Hill 100K.

New and unpublished LA-ICP-MS U-Pb detrital zircon ages from the Chapel Hill 100K includes: maximum depositional ages (MDA) of 533 Ma for sample CH100K-3675; MDA of 526 Ma for sample CH100K-3834; and MDA of 552 Ma for sample CH100K-Ben-165.

The previously existing and new age data indicates that volcanism and sedimentation of the Uwharrie Volcanic Complex spanned from ca. 552 Ma to ca. 526 Ma and was contemporaneous with the deposition of the Seagrove formation and deposition in the Albemarle Group (Figures 1 and 2).

ESTABLISHMENT OF THE SEAGROVE FORMATION (informal name)

New mapping in the Chapel Hill and Southern Pines 100Ks, where the Albemarle arc overlaps the Hyco arc, has confirmed the interpretation by Harris (1984) of the presence of a coarse-grained (sand-rich) submarine fan system that is contemporaneous with volcanism. Harris and Glover (1988) assigned their submarine fan deposits to the Aaron Formation. While Harris (1984) links the submarine fan to the Hyco arc, the new mapping of this study, coupled with new crystallization and detrital zircon ages, links the deposits to the Albemarle arc. This study establishes the Seagrove formation and defines its source as the volcanic rocks of the Uwharrie Volcanic Complex with a smaller component from the Hyco arc.

Extensive areas identified as Aaron Formation by Harris (1984) and Harris and Glover (1988) have been tentatively assigned to the newly defined Seagrove formation (this map). Additionally, some areas previously mapped as Uwharrie Formation are included in the Seagrove formation. More work is needed on the nature of the Aaron Formation; if the Aaron Formation turns out to be wholly Albemarle arc related, the Seagrove formation nomenclature can be abandoned.




Figure 1: Generalized stratigraphy of the Uwharrie Volcanic Complex and Seagrove formation.




Figure 2: Sketch of proposed redefined Uwharrie Volcanic Complex and newly defined Seagrove formation and their relationship to the Albemarle Group - ca. 560 to 523 Ma.

SEAGROVE FORMATION (informal name)

An overall defining characteristic of the Seagrove formation is the presence of quartz-bearing conglomerates, conglomeratic sandstones and sandstones interlayered with varying amounts of finer-grained volcanoclastic sediments that include bedded siltstones and mudstones. Individual units of the Seagrove formation are generally defined by the abundance of coarse-grained clastic sediments with respect to fine-grained clastic sediments. Proximal facies (proximal to Uwharrie Volcanic Complex units) are dominated by immature tuffaceous conglomerates, conglomeratic sandstones and sandstones with abundant clasts of rhyodacite and crystal fragments of plagioclase and quartz. Quartz crystal content varies locally. Medial to distal facies are dominated by conglomeratic siltstones (graywacke), sandy siltstones, siltstones, bedded siltstones and mudstones. Lenses of quartz-bearing conglomeratic sandstones and sandstones are locally present (Figure 3).

Data from Seiders's (1981) map in the Asheboro area was utilized in the compilation effort for the Chapel Hill 100K and for on-strike units in the Southern Pines 100K. Following accepted regional stratigraphy, Seiders (1981) separated his units from oldest to youngest, into the Uwharrie Formation and the Tillery and Cid Formations of the Albemarle Group (Stromquist and Sundelius, 1969 and Milton, 1984). The units of Seiders (1981) were maintained for the most part except in areas where new or recent age dates required modifications (ex. portions of the unit CZufc (felsic volcanoclastic rocks) on the east side of the Seiders (1981) map were reassigned to the Seagrove formation and CZups (porphyritic felsite near Seagrove) was reassigned to the Morrow Mountain Rhyodacite).

SEAGROVE FORMATION NEW GEOCHRONOLOGY DATA

Pollock et al. (2010) reported a youngest detrital zircon of ca. 545 Ma from the Erect member of the Uwharrie Formation (now included in the Seagrove formation). Recent LA-ICP-MS U-Pb zircon analyses of sandstones from Seagrove formation units in the Chapel Hill 100K indicate the presence of abundant detrital zircons associated with the Albemarle arc (Bradley, 2023; Pelt and Bradley, 2023). A re-evaluation of these zircon analyses yields the following: for NCGS sample Coleridge-3055, there is a distinct age distribution that is < 600 Ma and all MDAs are between ca. 545 Ma and ca. 549 Ma; for NCGS sample Bear Creek-626, all MDAs are between ca. 551 Ma and ca. 560 Ma (T. LaMaskin, personal communication, 2025).

New and unpublished LA-ICP-MS U-Pb zircon analyses from additional Seagrove formation unit samples in the Chapel Hill and Southern Pines 100K map areas includes: MDA of 559 Ma for sample COL-8005 in the Chapel Hill 100K; MDA of 554 Ma for sample COL-472 in the Chapel Hill 100K; MDA of 543 Ma for sample SP100K-550363 in the Southern Pines 100K; MDA of 533 Ma for sample CH100K-3675 in the Chapel Hill 100K; MDA of 526 Ma for sample CH100K-3834 in the Chapel Hill 100K; MDA of 524 Ma for sample SP100K-550946 in the Southern Pines 100K; and MDA of 523 Ma for sample CH100K-3839 in the Chapel Hill 100K. A summary of existing and new age dates within the Southern Pines 100K map are provided in Table 1. For a summary of existing and new age dates in the Chapel Hill 100K, see the Open-file report of the Chapel Hill 100K (Bradley et al., 2025a).

The previously existing and new age data indicate that sedimentation of the Seagrove formation spanned from ca. 560 Ma to ca. 523 Ma and was contemporaneous with the Uwharrie Volcanic Complex and deposition in the Albemarle Group.




Figure 3: Preliminary interpretation of the environment of deposition of the newly defined Seagrove formation with respect to the Uwharrie Volcanic Complex. Base figure after Stow and Mayall (2000) and Reading and Richards (1994) of a sand-rich submarine fan system. Green units are andesitic to basaltic volcanics.

ALBEMARLE GROUP

TILLERY FORMATION

Tillery Formation mudstones (CZtm) and intermediate to mafic volcanics (CZtimv) are present in the northwest corner and west-central portion of the map area. These map areas are immediately on strike with mapped units in the Charlotte 250K (Goldsmith et al., 1988) and the more detailed maps of Stromquist and Henderson (1985) and Conley (1962a). Detailed mapping by Upchurch (1968) interpreted the contact of the Tillery Formation (argillite facies unit of Upchurch (1968)) and units to the east (tuffaceous-sediment facies of Upchurch (1968) now mapped as part of the Seagrove formation) as gradational. Mapping of this study interprets the contact between the Tillery Formation and Seagrove formation as gradational.

ALBEMARLE ARC RELATED PLUTONIC ROCKS

Three small outcrop areas of fine-to medium-crystalline granodiorite (Zagd-p), interpreted to be related to the Parks Crossroads pluton, are present in the map area. Two of the outcrop areas are located northwest of Ether. One is surrounded by the CZsehc unit, the other is surrounded by unit CZmmr-ps. The third, and largest of the outcrop areas, is located southwest of Dover and is surrounded by unit CZseb.

The main outcrop area of the Parks Crossroads pluton (Zagd-p) is located in the west-central portion of the Chapel Hill 100K. Morrison and Coleman (2023) reported a new U-Pb zircon age for the Parks Crossroads pluton of ca. 552 Ma. Metamorphosed volcanoclastic rocks of the Seagrove formation have MDAs between ca. 551 Ma and ca. 560 Ma (T. LaMaskin, personal communication, 2025). As such, portions of the Seagrove formation units may have been disconformably deposited on top of the exhumed and eroded plutonic rocks. Unit CZmmr-ps has a crystallization age of ca. 536 Ma (Bradley et al., 2025); as such, the outcrop area is likely a xenolith of older rock in the younger rhyodacite.

ELLERBE AREA VOLCANICS AND VOLCANICLASTIC ROCKS

West of the Town of Ellerbe is a section of mainly fine-grained metamorphosed volcanosedimentary rocks consisting of bedded to laminated siltstones and mudstones with intermediate to mafic lavas and volcanoclastic rocks locally present. This area is separated from the main section of the Carolina terrane by the Wadesboro sub-basin of the Deep River Triassic basin. The units are interpreted to extend to the south, to the City of Rockingham, and are covered by Coastal Plain sediments with outcrops occurring only in the larger streams. These units are separated from the Easternmost Carolina terrane to the east by a possible ductile normal fault similar to the Fishing Creek fault in the Henderson 100K (Blake et al., 2025 and Blake et al., 2012).

The rocks are separated into three units that include: 1) metamorphosed laminated mudstones (CZelmd), 2) metamorphosed thin- to thickly-bedded mudstones (CZemd), and 3) metamorphosed intermediate to mafic volcanics and clastic rocks (CZeimvc). The laminated mudstones are interpreted to be the oldest unit and have visual similarities with the Tillery Formation. The thin- to thickly-bedded mudstones and intermediate to mafic volcanics and clastic rocks have lithologic and visual similarities with the Cid Formation.

New and unpublished LA-ICP-MS U-Pb detrital zircon ages from unit CZemd include: maximum depositional ages (MDA) of 540 Ma for sample SP100K 551983 and MDA of 562 Ma for sample SP100K 551641. A new and unpublished LA-ICP-MS U-Pb detrital zircon age from unit CZelmd yields a maximum depositional age (MDA) of 543 Ma for sample SP100K 551786.

FOLDS IN THE CAROLINA TERRANE

Folds in Hyco and Aaron Formations

The Hyco Arc and Aaron Formation lithologies were folded and subjected to low grade metamorphism during the ca. 578 to 554 Ma (Pollock, 2007; Pollock et al., 2010) Virgilina deformation (Glover and Sinha, 1973; Harris and Glover, 1985; Harris and Glover, 1988; and Hibbard and Samson, 1995). In the Chapel Hill 100K, original layering of Hyco Formation lithologies is observed, ranging from shallowly to steeply dipping. The range of structural attitude is interpreted to be a result of open to tight folds that are locally overturned.

Rocks in the map area were also subjected to the ca. 450 Ma Cherokee deformation and low-grade metamorphism (Hibbard et al., 2010 and 2012). Evidence of Cherokee deformation in the map area is scarcely seen in Hyco Formation units. Evidence is best exposed in area pyrophyllite mines where outcrop-scale folds deform an earlier foliation (see field trip stop 7 in Clark et al., 2011). This folding is associated with local deformation along several identified high-angle reverse faults in the map area (e.g. Glendon Fault and Robbins Fault).

In general, Hyco Formation rock units have been affected by the Virgilina and Cherokee deformations. Both deformations were generally co-planar, and as such, the metamorphic foliations in most locations are indistinguishable. The typical close-to-tight fold geometry of Hyco related units may be the result of the tightening of Virgilina deformation folds by the Cherokee deformation.

Folds in Albemarle arc units

Albemarle arc units have been affected by the Cherokee deformation. Folds attributed to the Cherokee deformation are generally open folds and include the well-known Silver Valley Syncline, Denton Anticline, New London Syncline, and Troy Anticline in the western portions of the terrane. The hinge line of the Troy Anticline enters the map area near the west-central edge of the map and extends toward the northeast and ends within a Hyco Formation unit interpreted as a local culmination of the Troy Anticline. The anticline then enters a depression, extending to the northeast where it exposes additional Hyco Formation rocks. The southeastern limb of the Troy Anticline is disrupted by high-angle reverse and thrust faulting. The Peachland syncline (Randazzo, 1972 and Moye, 2018) is a parasitic fold located south of the Troy anticline, which plunges southwest and exposes Tillery Formation metasediments to the south of Wadesville. The hinge of the Peachland syncline is traced discontinuously northeast for a few miles.

FAULTS

High-angle reverse faults

Several terrane-internal ductile faults have been recognized in Chatham, Moore and Montgomery counties within the Carolina terrane. The best known are the Glendon and Robbins faults, which have long been recognized by past workers (e.g. Stuckey, 1928 and Conley, 1962b). The Glendon Fault is a high angle reverse fault that is discontinuous and a locus of pyrophyllite alteration for multiple miles from northeast Moore County into southern Chatham County. Portions of the Glendon Fault are interpreted to be parallel to the axial surfaces of regional-scale overturned folds and locally disrupting an anticline near its crest (Green et al., 1982 and Klein, 1985). In general, the Glendon Fault is a zone of intense deformation ranging from 10 to 50 meters wide with abundant small-scale folds and fractures along with deformed and undeformed quartz veins, indicating a complicated movement history (Klein, 1985). Quartz veins may be folded and high-strain foliations are present within the fault zone, overprinting and/or transposing primary bedding and regional foliation. The Robbins Fault, present in the Robbins area, is well documented by Powers (1993), and like the Glendon Fault, is a zone of pyrophyllite alteration and complex deformation. Main movement on the Glendon and Robbins faults is speculated to be related to the Cherokee deformation.

Several other high-angle reverse faults with varying degrees of hydrothermal alteration were identified during mapping. In Chatham County, metamorphic foliation data indicates that the dip of the foliation progressively becomes shallower to the southeast approaching the high-angle faults. In the immediate area of the faults, sericite ($\pm$ pyrophyllite) phyllites and schistose phyllites with composite-like fabrics are common. It is interpreted that the older foliation has been transposed to a younger phyllonitic foliation within the fault zones.

Thrust faulting in the Carolina terrane?

Newly collected structural data by NCGS staff and compiled structural data from Carpenter (1999) and Hibbard (unpublished field data) in the Erect 7.5-Minute Quadrangle area of the Chapel Hill 100K, indicate abundant anomalous shallow dipping foliations and primary beddings consistently dipping northwest. The foliations typically dip less than 54 degrees to the northwest with a significant population of dips between 18-36 degrees. Mineral lineation data is sparse but indicates possible stretching in a northwest to southeast parallel orientation. The nature of these low-lying foliations is not well understood but may be related to thrust faulting and the presence of thrust duplexes(?). Tentatively, a thrust fault has been mapped separating hydrothermally altered volcaniclastic rocks of the Hyco Formation and Seagrove formation rock types. If it's an accurate interpretation, other un-mapped thrust faults may be present in the surrounding areas.

Eastern Piedmont Fault System

In the eastern portion of the map are strands of the Eastern Piedmont Fault System (EPFS), which are mostly buried by Coastal Plain sediments (Hatcher et al., 1977; Bobyarchick, 1981 and Stoddard et al., 1991). The EPFS is a system of Late Paleozoic-aged right-lateral ductile faults that are northeast-trending and mark terrane boundaries. The Nutbush Creek Fault and Neuse River Fault zones are mapped as continuations from exposed areas to the northeast of the Southern Pines 100K. The Barbeque fault is a

newly identified inferred fault that separates the Easternmost Carolina terrane from the newly identified amphibolite facies Upper Little River terrane.

Other faulting

Abundant evidence of brittle faulting is present at the outcrop-scale and map-scale as well as large-scale lineaments (as interpreted from hillshade LiDAR data). Brittle faults and lineaments are interpreted to be associated with arc rifting (Moye, 2023b) and Mesozoic extension. Numerous northwest-southeast trending brittle faults are mapped within the Carolina terrane portion of the map. It is speculated that these faults may have formed during arc-rifting in the Neoproterozoic to Paleozoic and were later reactivated during the Mesozoic. Within the Triassic basin of the map area, there are several named major Mesozoic brittle normal faults, which include the: Jonesboro Fault, Governors Creek Fault, and Crawley Creek Fault. Numerous map-scale relay ramps have been mapped by Reinemund (1955) on the west and east sides of the basin.

LiDAR LINEAMENTS

Linear geomorphic features interpreted from hillshade LiDAR are included on the map. The inclusion of LiDAR lineaments is an outgrowth from an NCGS derivative publication – *Groundwater Features Map of Chatham County and Surrounding Areas* (Bradley and Bolich, 2022).

The hillshade LiDAR dataset displays a visually striking network of lineaments that are parallel to diabase dikes and known orientations of brittle faults. Some of these lineaments can be traced for multiple miles across the map area and have linear topographic expressions that are 100's to greater-than 500 feet wide in some locations. The lineaments are interpreted to be fracture zones; however, field observations could not confirm the presence of each potential brittle fault. The map area has hundreds of lineaments when viewed at varying scales. Only the major lineaments that are easily visible at 1:50,000- to 1:100,000-scale are identified on the map.

Multiple unmarked geomorphic lineaments are visible when viewed at 1:24,000- to 1:50,000-scale on the elevation data from LiDAR. These minor lineaments extend for short distances (100's to 1000's of feet). Minor lineaments are not indicated on the map but can be identified by a close inspection of the LiDAR data. These minor lineaments may have important implications for site-specific investigations. Site specific studies should review lineaments at the 1:24,000-scale or greater.

EASTERNMOST CAROLINA TERRANE

The Easternmost Carolina terrane is located east of the Jonesboro Fault and west of the Upper Little River terrane. In the map area, the terrane is underlain by metamorphosed crystalline rocks of the Cary sequence (Parker, 1979; Farrar, 1985) and meta-intrusive units. The Cary sequence is interpreted to be part of the Carolina terrane but separated from the rest of the terrane by the Triassic basin (Hibbard et al., 2002). One of the main rock units is the Big Lake-Raven Rock schist. In the Cary Quadrangle in the Raleigh 100K, a sample from the unit yielded discordant $^{207}\text{Pb}/^{206}\text{Pb}$ zircons ages of 573, 574, and 579

Ma, and an upper intercept age of 575 ± 12 Ma, interpreted as the time of crystallization (Goldberg, 1994).

The terrane also includes metamorphosed intrusive rocks that are interpreted to be subvolcanic feeders to the volcanic rocks of the Cary Sequence (Blake et al., 2001). The Lemon Springs pluton is an example of metamorphosed intrusive rocks present in the terrane. A Rb-Sr date of 429 ± 10 Ma (Kish and Campbell, 1986) for the Lemon Springs pluton likely indicates a minimum age.

UPPER LITTLE RIVER TERRANE

Compilation work using unpublished data from Robert J. Butler and new data collected by NCGS staff have identified a map scale body of schist with garnet porphyroblasts up to 4 mm in diameter. The map unit is interpreted to be amphibolite facies and separated from the Easternmost Carolina terrane by a ductile fault. Large outcrops are exposed on Upper Little River. Additional work is needed to confirm the presence of a ductile fault (tentatively named Barbeque Creek fault(?) after the nearby Barbeque Creek and the community of Barbeque) along the western boundary of the terrane. To the east, the terrane is bounded by the Nutbush Creek Fault. The unit extends into the Fayetteville 100K.

CRABTREE TERRANE

The Crabtree terrane is present in the subsurface but covered by Coastal Plain Sedimentary deposits. Generally, the Crabtree terrane includes amphibolite facies schists and phyllites. The rock unit in the map area is interpreted as sericite phyllite using geophysical and boring data from Lawrence and Hoffman (1993). The Crabtree terrane is bounded on the west by the Nutbush Creek Fault. The Neuse River Fault separates the Crabtree terrane from the Spring Hope terrane to the east.

SPRING HOPE TERRANE

The Spring Hope terrane is present in the subsurface but covered by Coastal Plain Sedimentary deposits. Generally, the Spring Hope terrane includes metamorphosed sedimentary and volcano-sedimentary rocks. The rock units in the map area were interpreted using geophysical and boring data from Lawrence and Hoffman (1993). The Spring Hope terrane is separated from the Crabtree terrane by the Neuse River Fault.

PERMIAN TO MISSISSIPPIAN INTRUSIVE ROCKS

The Lilesville Granite (Waskom and Butler, 1971 and Waskom, 1970) is a distinctive potassium-feldspar megacrystic granite. The pluton is ring-shaped, intruded by the Pee Dee gabbro (located in the Laurinburg 100K) and has an exterior and interior contact aureole that is well documented by Evans and Speer (1984). The Millstone Lake Pluton is a small body that is mostly covered by Coastal Plain sediments. Abundant outcrops are present in Millstone Lake 4-H camp and have been quarried for millstones.

The Pinebluff pluton is a newly identified pluton that is completely covered by Coastal Plain sediments. Bartlet (1967) reported that a well driller encountered coarse-crystalline granite in the Town of Pinebluff at the former “Aberdeen Ice House” at a depth of approximately 100 feet. Additionally, NCDOT borings at bridge No. 14 on SR1172 reported granite at a depth of approximately 40 ft. The pluton is present as a distinct circular body on a preliminary magnetic survey map from the Piedmont Bridge South Magnetic and Radiometric Survey (USGS Piedmont Bridge, 2025).

CONTACT AUREOLE OF THE LILESVILLE GRANITE

The contact aureole of the Lilesville Granite is well documented by Evans and Speer (1984). Regional metamorphosed greenschist-facies sediments of the Carolina terrane have been overprinted by contact metamorphism that produced distinct hornfels and migmatitic gneisses. This mapping separates the contact aureole following Evans and Speer (1984) into three units: 1) exterior aureole hornfels, 2) interior aureole hornfels, and 3) migmatized interior aureole hornfels. The only contact aureole unit present in the Southern Pines 100K is the exterior aureole hornfels.

DEEP RIVER TRIASSIC BASIN

A significant portion of the central area of the map is underlain by Triassic-aged sedimentary rocks of the Deep River Triassic basin. A review of the basin geology is provided in Clark et al. (2001). The basin is separated into three sub-basins: Durham, Sanford and Wadesboro. The Colon cross-structure (Campbell and Kimball, 1923, and Reinemund, 1955), located on the Chapel Hill 100K map, is a constriction zone in the basin characterized by crystalline rocks overprinted by complex brittle faulting. The Colon cross-structure marks the transition between the Durham and Sanford sub-basins. The Sanford and Wadesboro sub-basins are present on the Southern Pines 100K map and are interpreted to be continuous but separated by the Pekin cross-structure and under Coastal Plain sedimentary deposits (Olsen et al., 1991). The Ellerbe basin, located in the southern portion of the Southern Pines 100K, is separated from the Wadesboro sub-basin by crystalline rock and Coastal Plain sedimentary deposits. The Ellerbe basin is probably the remnant of Triassic basin fill on top of an exhumed rider block system (Olsen et al., 1991). The Jonesboro Fault is interpreted to bound the Ellerbe basin to the east.

SANFORD AND WADESBORO SUB-BASINS AND ELLERBE BASIN MAP UNITS

In the Sanford sub-basin, three stratigraphic units have been identified and formalized, from oldest to youngest, as the Pekin, Cumnock and Sanford Formations (Campbell and Kimball, 1923 and Reinemund, 1955). A majority of the geologic contacts within the Sanford sub-basin in the northern portion of the Southern Pines 100K were digitized as presented by Reinemund (1955).

This compilation work included the digitization of existing maps of Randazzo and Copeland (1976) and Brazell (2013) in the Wadesboro sub-basin and mapping by Dineen (1982) in the small Ellerbe basin. Contacts were modified if newly collected field data indicated a need for revisions. Two stratigraphic units are identified in the Wadesboro sub-basin, which include the Pekin Formation and the newly

defined Wadesboro formation (Brazell, 2013). The community of Pekin is located within the Wadesboro sub-basin and is the type locality of the Pekin Formation (Cambell and Kimball, 1923).

The Ellerbe basin was mapped by Dineen (1982) and includes sedimentary breccias, conglomerates, sandstones and mudstones that are not formally linked to known formations.

The contact between the Sanford and Wadesboro sub-basins was chosen at a significant aeromagnetic linear feature that may be a fault intruded by a diabase dike. This contact is for cartographic purposes to separate the sub-basins. The contact location is buried by Coastal Plain sediments and was chosen to demarcate the sub-basins in the vicinity of the Pekin Cross structure. The Sanford Formation in the northern portion of the map area was terminated under Coastal Plain sediments at a significant aeromagnetic linear feature. Triassic sediments southwest of the Sanford Formation are assigned to an undifferentiated unit (Tru). The limited exposure in this area and fine-grained nature of the sediments (not indicative of the Sanford Formation) was not sufficient to assign the rock to existing Formations.

POST-TRIASSIC ROCK UNITS and COASTAL PLAIN SEDIMENTARY DEPOSITS

Dikes of Jurassic diabase intrude the Triassic sedimentary and older crystalline rocks of the map area. Coastal Plain Sedimentary deposits cover a majority of the bedrock in the Southern Pines 100K. The general distribution of Coastal Plain deposits is provided on an inset figure on the map. Thin residual Coastal Plain deposits are common in locations away from the main outcrop area. In the extreme southeast of the quadrangle, Coastal Plain deposits may be as thick as 240 feet (well data from CD-P-1-67 reported in Lawrence and Hoffman (1993)).

Quaternary alluvium is present in most modern river valleys, with at least two levels of fluvial terraces along the major drainages. These terraces, where preserved, likely mark the location and elevation of ancestral river systems, prior to incision to the modern floodplain levels (Grimley, 2023).

ACKNOWLEDGEMENTS

This geologic mapping was funded in part by the USGS National Cooperative Geologic Mapping Program under STATEMAP, award numbers G24AC00339, 2024 and G25AC00477, 2025.

The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

Thanks to Jim Hibbard for the consultation and input on stratigraphy of the Carolina terrane.

Thanks to David E. Blake for consultation on extent of Eastern Piedmont Fault system and terranes beneath Coastal Plain sediments.

DESCRIPTION OF MAP UNITS

LATEST PERMIAN TO EARLY CENOZOIC ROCKS

FAULT ROCKS

CzMzqv2 **Vein quartz and quartz breccia (Cenozoic-Mesozoic)**—White to light gray, coarse-crystalline, highly fractured and cataclastic vein quartz. Locally includes subhedral quartz crystals as vein infilling.

INTRUSIVE ROCKS (UNMETAMORPHOSED)

Jd **Diabase (Jurassic)**—Black to greenish-black, fine- to medium-grained, dense, consists primarily of plagioclase, augite and may contain olivine. Locally has gabbroic texture. Occurs as dikes up to 100 ft wide. Diabase typically occurs as spheriodally weathered boulders with a grayish-brown weathering rind. Red station location indicates outcrop or boulders of diabase. Purple station locations indicate outcrop or boulders of gabbroic textured diabase.

MZlamp **Lamprophyre (?) (Mesozoic to Neoproterozoic)**—Gray to pinkish gray, fine- to medium-grained, exceptionally dense, with alkali-feldspar, plagioclase and amphibole. The groundmass consists of alkali feldspar and plagioclase, with alkali feldspar more abundant than plagioclase. Plagioclase crystals (>1 cm) also occur are subhedral and commonly zoned. Locally, amphibole occurs in elongate slender prismatic habit (1-4 mm) and is randomly oriented. Sparse amygdules of quartz(?) up to 5 mm present locally. The rock is unmetamorphosed but may have magmatic and/or hydrothermal alteration. Occurs as dikes that are

coincident with diabase. Outcrop and boulders are typically spheriodally weathered with reddish-brown weathering rind. Red square station locations mark outcrops or boulders.

TRIASSIC SEDIMENTARY ROCKS OF THE NEWARK SUPERGROUP

DEEP RIVER BASIN

TrU Undifferentiated Triassic sediments in the Deep River basin (Triassic)—

Brown to maroon, locally red, interbedded mudstones, siltstones and sandstones. Majority of area is covered by Coastal Plain sedimentary deposits. Exposure is rare and present only along Naked, Drowning and Jackson creeks. Insufficient research has been done to confidently assign to a named formation. May be in distinct structural block(?) bounded to west by crystalline rock high and to east by a significant diabase dike identified on aeromagnetic data. The diabase may intrude a major fault separating distinct structural blocks in the basin(?).

SANFORD SUB-BASIN

Trs Sanford Formation (Triassic)—Mainly red to brown, locally purple, coarse-grained, arkosic sandstones and conglomerates. Subordinate amounts of claystone, siltstone and fine-grained sandstone (Reinemund, 1955).

Trsc Conglomerate of the Sanford Formation (Triassic)—Mainly conglomerates with fragments of metamorphic rock and quartz embedded in and interbedded with red mudstone, siltstone and sandstone (Reinemund, 1955). Equivalent to Trcc of Lithofacies Association III in the Cokesbury Quadrangle.

Trc **Cumnock Formation (Triassic)**—Gray and black claystone, shale and siltstone.

Gray sandstone. Contains beds of coal and carbonaceous (organic-rich) shale (Reinemund, 1955). Includes coal horizons.

Trp **Pekin Formation (Triassic)**—Gray, Brown to maroon, white mica bearing, interbedded mudstones, siltstones and arkosic sandstones. Outcrops and boulders of float identified as part of Pekin Formation are strongly indurated compared to conglomerates identified as part of Chatham Group. Identified as the Pekin Formation by Reinemund (1955).

Trpc **Conglomerate of the Pekin Formation (Triassic)**—Reddish-brown to dark brown to purplish-red, irregularly bedded, poorly sorted, cobble to boulder conglomerate. Clasts are chiefly miscellaneous felsic and intermediate metavolcanic rocks and quartz. Typically present adjacent to border faults. Outcrops and boulders of float identified as part of Pekin Formation are strongly indurated compared to conglomerates identified as part of Chatham Group. Identified as the Pekin Formation-basal conglomerate by Reinemund (1955).

Trpwpp **Zone of wind polished and faceted pebbles in the Pekin Formation (Triassic)**—Area in Pekin Formation with wind polished and faceted pebbles. Interpreted by Reinemund (1955) as an indication of environment of deposition in which pebbles exposed on the surface during dry seasons were wind polished and faceted.

WADESBORO SUB-BASIN

Trw **Wadesboro Formation (Triassic)**—Thin (1 to 5 meters thick), sheet siltstones and sandstones interbedded with thin beds of mudstone and siltstone.

Sandstone to mudstone ratio of about 1:3. Local occurrence of conglomeratic interlayers present mainly proximal to faults. The sandstones of the Wadesboro Formation are more quartz rich (compared to Pekin) and are generally classified as feldspathic litharenites and litharenites. Lithic components of the sandstones are generally more sub-angular to sub-rounded than Pekin Formation and are dominated by plutonic and metamorphic rock fragments. Described by Brazell (2013).

Trp-w **Pekin Formation of the Wadesboro sub-basin (Triassic)**—Thick (greater than 5 meters thick) channelized deposits that contain trough cross-bedded coarse to medium grained sandstones interbedded with thick deposits of mudstone, siltstone, and shale that preserve paleosols. The sandstone to mudstone ratio is about 1:5. Grains in the sandstones of the Pekin Formation are generally angular to sub-angular and have a large component of feldspars, sedimentary lithics, and clay to silt sized particles filling the pore spaces between larger grains. Described by Brazell (2013).

Trpc-w **Conglomerate of the Pekin Formation of the Wadesboro sub-basin (Triassic)**—Poorly sorted, cobble to boulder conglomerate interlayered with conglomeratic sandstone, sandstone and siltstone. Clasts are chiefly subrounded to rounded miscellaneous felsic and intermediate metavolcanic rocks and quartz. Typically present adjacent to border faults.

Trgb-w **Granitic breccia of the Wadesboro sub-basin (Triassic)**—Gravel- to cobble-

sized, sedimentary breccias composed of granitic clasts. Identical in texture and composition as unit Tregb in the Ellerbee basin.

ELLERBE BASIN

- Trem **Mudstone of the Ellerbe Triassic basin (Triassic)**—Red to dark reddish brown with red and gray mottling, mudstone composed of quartz, clays, mica and fragments of Carolina terrane siltstones and granite. Beds range from less than 1 foot to more than 30 feet thick. Described by Dineen (1982).
- Tres **Sandstone of the Ellerbe Triassic basin (Triassic)**—Dark red to pale red, fine - to coarse-grained, poorly to moderately sorted, range from structureless to thinly planar laminated, clayey quartz and arkosic sandstones. Bed thickness commonly less than 3 feet but locally up to 10 feet. Described by Dineen (1982).
- Treb **Breccia of the Ellerbe Triassic basin (Triassic)**—Reddish brown to dark reddish brown, gravel- to boulder-sized, sedimentary breccias composed of clasts of Carolina terrane bedded to laminated mudstones and siltstones (argillites), vein quartz, granite and occasional Triassic sandstone and mudstone rip-up clasts (Dineen, 1982). Breccias grade basinward into conglomerate composed of rounded clasts.
- Tregb **Granitic breccia of the Ellerbe Triassic basin (Triassic)**—Gravel- to cobble-sized, sedimentary breccias composed of granitic clasts (Dineen, 1982). In contact with the Lilesville granite.

PERMIAN TO MISSISSIPPIAN INTRUSIVE ROCKS

- PPMlg **Lillesville Granite (Permian-Mississippian)**—Coarse-crystalline, hypidiomorphic inequigranular, biotite granite and granodiorite with alkali feldspar megacrysts up to 4 cm long. Speer (1988) describes the mineralogy as follows: the alkali feldspar is microcline microperthite that occurs as pink, anhedral to subhedral crystals less than 2 cm long and blocky subhedral megacrysts up to 4 cm across. White to greenish plagioclase occurs as smaller subhedral grains with normal oscillatory zoning (An30 to An80) and local antiperthite. Biotite occurs as flakes up to 4 mm across and is pleochroic in shades of brown. Quartz is anhedral, gray and less than 5 mm. Accessory minerals include allanite, zircon, apatite and opaques. Secondary minerals include muscovite, chlorite, epidote and carbonate. Fullagar and Butler (1979) report a Rb-Sr date of 326 ± 27 Ma.
- PPMmlp **Millstone Lake Pluton (Permian-Mississippian)**—Pink-white to pink-gray, medium- to coarse-crystalline, biotite granite. Outcrops in creeks near Millstone 4-H Camp. Used for making millstones by early settlers. Extent unknown with majority of pluton covered by Coastal Plain sediments. Named Millstone Lake Pluton in McSween et al. (1991).
- PPMpbb **Pinebluff pluton (Permian-Mississippian)**—Dark gray to gray, coarse-crystalline granite in the subsurface under Coastal Plain sediments. Reported in Bartlett (1967) and in a NCDOT boring. Distinct circular body present on Piedmont Bridge South Magnetic and Radiometric Survey (USGS Piedmont Bridge, 2025).

SPRING HOPE TERRANE

METAVOLCANIC AND METASEDIMENTARY UNITS

- CZfv-gp1 **Metamorphosed felsic volcanic rocks interpreted from geophysical methods 1 (Cambrian-Neoproterozoic)**—Area in subsurface interpreted as felsic volcanic rocks using geophysical data from Lawrence and Hoffman (1993). Covered by Coastal Plain sedimentary deposits.
- CZmsd-gpb **Metamorphosed sediments interpreted by geophysical methods and boring data (Cambrian-Neoproterozoic)**—Area in subsurface interpreted as sericite phyllite and metamudstone in the northeast portion of the unit in the Fayetteville 100K and metamorphosed graywacke and mudstone in the southwest portion of the unit in the western portions of the Fayetteville 100K and eastern portions of the Southern Pines 100K using geophysical and boring data from Lawrence and Hoffman (1993). Metamorphosed conglomerate, sandstone and siltstone are present at Stevens Mill west of Goldsboro in the Fayetteville 100K. Covered by Coastal Plain sedimentary deposits.
- CZmsd-gpb2 **Metamorphosed sediments interpreted by geophysical methods and boring data 2 (Cambrian-Neoproterozoic)**—Area in subsurface interpreted as phyllitic siltstone and slightly metamorphosed mudstone using geophysical and boring data from Lawrence and Hoffman (1993). Covered by Coastal Plain sedimentary deposits.
- CZmsd-gpb3 **Metamorphosed sediments interpreted by geophysical methods and boring data 3 (Cambrian-Neoproterozoic)**—Area in subsurface

interpreted as slaty metamudstone using geophysical and boring data from Lawrence and Hoffman (1993). Covered by Coastal Plain sedimentary deposits.

CRABTREE TERRANE

METAVOLCANIC AND METASEDIMENTARY UNITS

CZsph-gpb **Sericite phyllite interpreted from geophysical methods and boring data**

(Cambrian to Neoproterozoic)—Area in subsurface interpreted as sericite phyllite using geophysical and boring data from Lawrence and Hoffman (1993). Located in the Crabtree terrane (?) in between the Nutbush Creek and Neuse River faults. Covered by Coastal Plain sedimentary deposits.

CZcimv-gp **Intermediate to mafic volcanics in the Crabtree terrane interpreted from**

geophysical methods (Cambrian to Neoproterozoic)—Area in subsurface interpreted as Intermediate to mafic volcanic rocks using geophysical data from Lawrence and Hoffman (1993). Located in the Crabtree terrane (?) in between the Nutbush Creek and Neuse River faults. Covered by Coastal Plain sedimentary deposits.

UPPER LITTLE RIVER TERRANE

CZulrs **Upper Little River schist (Cambrian to Neoproterozoic)**—Silvery-gray, fine - to medium- crystalline, garnet + white mica schist. Garnets porphyroblasts up to 4 mm. Large portions of unit covered by Coastal Plain sedimentary deposits.

EASTERNMOST CAROLINA TERRANE

METAINTRUSIVE UNITS

BUCKHORN DAM META-INTRUSIVE SUITE

CZbg **Meta-granitoid rocks of the Buckhorn Dam intrusive suite (Cambrian to Neoproterozoic)**—Dark-colored (CI=15-30), medium- to fine-grained, metatonalite, metagranodiorite and metagranite with variably developed foliation; composed mainly of plagioclase, quartz, epidote, microcline, biotite, and opaque minerals, with minor amounts of sericite, sphene, chlorite, and garnet. The more felsic granitoid rocks are mineralogically and chemically similar to the felsic metavolcanic rocks described below, and are probably the intrusive equivalents. The unit includes a number of small granitoid bodies, probably originally dikes and plugs, intruding felsic metavolcanic rocks northeast of the main outcrops of Buckhorn Dam intrusive suite.

CZbg-lsp **Lemon Springs pluton of the Buckhorn Dam intrusive suite (Cambrian to Neoproterozoic)**—Leucocratic to mesocratic, medium- to fine-crystalline, metamorphosed alkali granite with variably developed foliation. In hand-sample, mineral boundaries are not well defined except for those of tabular albite crystals and small (0.5 to 2.0 mm) pods of dark minerals that are acicular stilpnomelane (Campbell 1984). According to Campbell (1984), the average major mineral constituents from point counts are microcline (34%), quartz (31%), and albite (27%). Minor amounts of epidote, stilpnomelane, biotite, chlorite, white mica,

garnet, opaques, sphene, calcite, apatite, allanite, and pyrite are present, in order of decreasing abundance (Campbell, 1984). The Lemon Springs quarry also hosts, low-silica, high-potassium dikes that may represent metamorphosed lamprophyre and metamorphosed felsic and basaltic dikes (Campbell, 1984).

CZbg-rd **Rhyodacite to dacite of the Buckhorn Dam intrusive suite (Cambrian to Neoproterozoic)**—Gray, aphanitic, vitiric-like, flow-banded metamorphosed rhyodacite to dacite. May be an apophysis of the Lemon Springs pluton (?).

CARY METAMORPHIC SUITE

CZbr3 **Big Lake-Raven Rock schist 3 (Cambrian to Neoproterozoic)**—Light tan to orange-brown, fine- to medium-grained, white mica schist, phyllite and gneiss. Locally preserves primary volcanic texture, either fragmental or porphyritic. Inferred to have a dacitic volcanic and/or volcanoclastic protolith. Locally includes intermediate to mafic composition rocks that have been metamorphosed to mica phyllite.

CZbr-py **Phyllite and argillite (Cambrian to Neoproterozoic)**—Fine- to medium-crystalline phyllite and metamorphosed siltstones. Locally preserves primary sedimentary textures (bedding and crystal fragments). Tuffaceous interlayers present. May be a more distal and finer-grained facies related to Big Lake-Raven Rock schist units. Differentiated from unit to southwest (CZsphs-gpb) due to abundance of high magnetic signature bodies on Piedmont Bridge South Magnetic and Radiometric Survey (preliminary data, 2025). Large portions of unit covered by

Coastal Plain sedimentary deposits.

CZsphs-gpb

Sericite phyllite and metamorphosed sedimentary rocks interpreted by geophysical methods and boring data (Cambrian to

Neoproterozoic)—Area in subsurface interpreted as mainly sericite phyllite using geophysical and boring data from Lawrence and Hoffman (1993). Includes lesser amounts of fine-grained metamorphosed sedimentary rocks and metamorphosed granitoid. Northeast contact of unit is questionable and designates apparent contact between rocks of lower magnetic intensity to the southwest from rocks of higher magnetic intensity to the northeast. Outcrops in limited exposed areas include: phyllite, meta-rhyodacite, schistose phyllite and gneissic granitoid.

Majority of unit is covered by Coastal Plain sedimentary deposits.

CZmas

Mafic schist and phyllite and sericite phyllite (Cambrian to

Neoproterozoic)—Green, chlorite-actinolite schist, chlorite schist and sericite phyllite. Schist grade into dark green intermediate to mafic crystal tuffs. intercalated lithologies include epidosite, hematite rich epidote-quartzite breccia and chlorite phyllite. Small pyrite crystals are commonly found along the fine-grained quartz laminae. Hematite also commonly present. Small lenses of quartzo-feldspathic gneiss are intercalated with the unit. (Modified from Hicks, 1982)

CZimv-gp

Intermediate to mafic volcanics in the Easternmost Carolina terrane interpreted from geophysical methods (Cambrian to

Neoproterozoic)—Area in subsurface interpreted as Intermediate to mafic volcanic rocks using geophysical data from Lawrence and

Hoffman (1993). Located in the Easternmost Carolina terrane. Covered by Coastal Plain sedimentary deposits.

CAROLINA TERRANE

ALBEMARLE ARC

CZq **Quartz veins (Cambrian to Neoproterozoic)**—Area of abundant quartz present in a vein or set of veins. May include silicic alteration of country rock. Often vuggy texture present with terminating crystals.

ALBEMARLE ARC PLUTONS

CZagb-sm **Stony Mountain Gabbro (Cambrian to Neoproterozoic)**—Massive, grayish-green to dark-greenish-gray, fine to coarse grained. Composed of secondary amphibole, in part with cores of clinopyroxene, calcic plagioclase, epidote, chlorite, and quartz, with accessory sphene, apatite, calcite, phlogopite, and opaque minerals. Occurs as sheets, dikes, and irregular intrusive bodies; most common within well-bedded sedimentary and volcanic rocks. The sheet exposed west of Back Creek Lake has a basal zone rich in serpentine pseudomorphs after olivine; "comb layering" (Moore and Lockwood, 1973) is well exposed in the quarry at the top of the same sheet (Seiders and Wright, 1977). A sample collected from Ridges Mountain (west of Asheboro) in the unit yielded an age of ca. 545 Ma (DeDecker et al., 2013).

Zagd-p **Granodiorite of the Parks Crossroads pluton (Neoproterozoic)**—Leucocratic (Cl=5) , fine- to medium- grained, equigranular metamorphosed, granodiorite. Mineral assemblage includes quartz, plagioclase, and

green hornblende +/- chlorite, +/- epidote. Locally foliated in the Southern Pines 100K. A sample collected from an abandoned quarry in the unit in the Chapel Hill 100K yielded a weighted mean age of ca. 552 Ma (Morrison and Coleman, 2023). Contact with Hyco arc units interpreted as an intrusive contact. Contact with portions of the Seagrove formation units may be an unconformity based on preliminary detrital zircon data. Additional work is needed to confirm.

MORROW MOUNTAIN RHYODACITE SUITE

CZmmr-ps **Morrow Mountain rhyodacite – porphyritic felsite near Seagrove (Cambrian**

to Neoproterozoic)—Gray, rhyodacite (felsite of Seiders, 1981) with phenocrysts of albite and quartz. Occurs in a 3 km-wide belt and in small bodies on both sides of the belt. At the south end of the belt the felsite is flow layered, locally spherulitic, and the phenocrysts are relatively smaller and less abundant. To the north, in Randolph County, phenocrysts are larger (1-3 mm) and more abundant; flow layering and spherulites are generally absent. Both extrusive and shallow intrusive rocks may be represented. A sample collected from the southbound I-73/74 visitor center approximately 1.9 miles southwest of Seagrove in the Chapel Hill 100K yielded a LA-ICP-MS $^{206}\text{Pb}/^{238}\text{U}$ weighted means age of ca. 536 Ma (Bradley et al., 2025).

CZmmr-psa

Altered Morrow Mountain rhyodacite – porphyritic felsite near Seagrove (Cambrian to Neoproterozoic)—Light gray quartzose sericite phyllite; locally contains pyrite; commonly iron oxide-stained where weathered. Pyrophyllite locally present. Interpreted as hydrothermally

altered CZmmr-ps unit.

CZmmr-f **Morrow Mountain rhyodacite – felsite (Cambrian to Neoproterozoic)**—Light-gray to dark-gray rhyodacite (felsite of Seiders, 1981), aphanitic to porphyritic with phenocrysts of albite and quartz. Locally spherulitic and flow layered. May include lava flows, domes, necks, and dikes. Bradley et al. (2025), reported LA-ICP-MS 206Pb/238U weighted means age for a sample from Caraway Mountain in the Chapel Hill 100K of ca. 543 Ma. See table 1 for additional dates.

ALBEMARLE GROUP

TILLERY FORMATION

CZtm **Laminated to thin-bedded graded mudstone (Cambrian to Neoproterozoic)**—Typically grayish-green, less commonly gray, in beds 1 mm to 4 cm thick. Beds grade upward from lighter colored siltstone to darker claystone. Beds are extremely regular, laterally persistent, and show no crossbedding. Thick-bedded mudstone is uncommon. Rare beds of graded sandstone consist of volcanic rock and mineral fragments of diverse composition in an abundant argillaceous matrix. Contains biotite porphyroblasts

CZtfm **Felsite, felsic volcanoclastic rocks and interbedded, laminated- to thin-bedded mudstone (Cambrian to Neoproterozoic)**—Felsite, felsic volcanoclastic rocks, and interbedded, laminated- to thin-bedded mudstone

CZtimv **Intermediate to mafic volcanics in the Tillery Formation (Cambrian to**

Neoproterozoic)—Map scale bodies of metamorphosed gray-green, dark green and greenish-gray; typically unfoliated to weakly foliated, amygdaloidal, plagioclase porphyritic and aphanitic; andesitic to basaltic lavas and shallow intrusions. Hyaloclastic texture is common and imparts a fragmental texture on some outcrops and float boulders. Correlated with Ctb (basaltic rocks) of Stromquist and Henderson (1985).

CZtmc **Mafic volcanoclastic rocks (Cambrian to Neoproterozoic)**—Thin- to very thick bedded, grayish-green tuff, lapilli tuff, and tuff breccia. In addition to mafic volcanic debris, many rocks contain subordinate amounts of mudstone and felsic volcanic rock clasts

SEAGROVE FORMATION (INFORMAL)

CZsfc **Seagrove formation felsic volcanoclastic rocks (Cambrian to Neoproterozoic)**—Thin- to very thick bedded, light- to dark-gray, in part pale-green, tuff, lapilli tuff, and tuff breccia. The most abundant rock types are very thick bedded tuff, lapilli tuff, and tuff breccia, but in many parts of the map unit subordinate amounts of well-bedded, thin- to thick-bedded, in part graded, tuff, tuffaceous sandstones, tuffaceous conglomeratic sandstones and felsic volcanic mudstone are interbedded. In many rocks the principal clasts are dense felsite, like that occurring in the numerous felsite bodies of the formation. Clasts are angular to subrounded, very rarely well rounded. Other rocks have clasts with poorly preserved textures that are strongly suggestive of pumice. Such rocks occur throughout the unit but are especially

abundant in a 3 km wide belt, extending from south of Center Cross Church to Purgatory Mountain and located just west of the belt of porphyritic felsite at Seagrove (CZmmr-ps). These rocks generally lack interbeds of well-bedded tuff and may represent thick ashflow deposits. The probable pumice fragments are commonly much flattened but the flattening is in the plane of the regional cleavage or foliation and may be tectonic. Textural details are too poorly preserved to record most features that would clearly indicate welding.

CZsehc **Seagrove formation epiclastic rocks of the Harpers Crossroads area (Cambrian to Neoproterozoic)**—Metasedimentary package that ranges from fine-grained siltstones to coarse-grained sandstones, pebbly sandstones and conglomerates. The sandstones, pebbly sandstones and conglomerates commonly contain rounded to subangular clasts of quartz ranging from sand- to gravel-sized. In the sandstones, feldspar is the most prominent mineral grain; quartz varies from sparse to abundant in hand sample. Lithic clasts are typically prominent and range from sand- to gravel-size. Previously mapped as part of the Aaron Formation on open file maps of western Chatham County.

CZsehc-a **Altered Seagrove formation epiclastic rocks of the Harpers Crossroads area (Cambrian to Neoproterozoic)**—Area of Seagrove formation epiclastic rocks of the Harpers Crossroads area that is mildly to moderately hydrothermally altered. Alteration consists of silicified, sericitized and pyrophyllitized rock. Chloritoid locally present.

CZseb **Seagrove formation epiclastic rocks of the Biscoe area (Cambrian to**

Neoproterozoic)—Metasedimentary package that is dominantly fine- to medium-grained siltstones and sandy siltstones with local quartz bearing sandstones and pebbly sandstones. Lesser amounts of conglomerate. The sandstones and pebbly sandstones commonly contain rounded to subangular sand-size grains of quartz. Quartz grains appear to be less abundant than in CZsehc. In the sandstones, feldspar is the most prominent mineral grain; quartz varies from sparse to abundant in hand sample. When present, lithic clasts range from sand- to gravel-size and locally include pink-colored, angular clasts of altered (?) rhyodacite. Minor andesite to basalt lavas and/or shallow intrusions.

CZseb-a

Altered Seagrove formation epiclastic rocks of the Biscoe area

(Cambrian to Neoproterozoic)—Mixed unit of altered volcanoclastic sedimentary rocks within the CZseb unit. Alteration consists of silicified, sericitized and pyrophyllitized rock. Sulfides locally present. Vuggy quartz boulders and cobbles abundant throughout unit. Includes altered rhyodacite (CZmmr-f) that was likely emplaced as dikes.

CZsebimv

Seagrove formation epiclastic rocks of the Biscoe area with local

intermediate to mafic volcanics (Cambrian to Neoproterozoic)—Metasedimentary package that is dominantly fine- to medium-grained siltstones and sandy siltstones with local quartz bearing sandstones and pebbly sandstones identical to CZseb with local to abundant metamorphosed andesite to basalt lava and/or shallow intrusions. Local occurrence of metarhyodacite of CZmmr-f present.

CZsebimv-a

Altered Seagrove formation epiclastic rocks of the Biscoe area with

	local intermediate to mafic volcanics (Cambrian to Neoproterozoic)—Variably altered metasedimentary package that is dominantly fine- to medium-grained siltstones and sandy siltstones with local quartz bearing sandstones and pebbly sandstones with local to abundant metamorphosed andesite to basalt lava and/or shallow intrusions identical to CZsebimv. Includes phyllitic siltstone. Altered rock includes: sericite +/- pyrophyllite phyllite and abundant vuggy quartz veins. Includes areas of the historic Golconda, Uwarra, Iola and Martha Washington gold mines. Area is deeply weathered with only the most resistant rocks forming outcrops.
CZsimv	Seagrove formation intermediate to mafic volcanic rocks (Cambrian to Neoproterozoic)—Map scale bodies of metamorphosed light green, gray-green, gray, and dark gray; typically unfoliated to weakly foliated, amygdaloidal, plagioclase porphyritic and aphanitic; andesitic to basaltic lavas and shallow intrusions. Hyaloclastic texture is common and imparts a fragmental texture on some outcrops and float boulders.
CZsimvc	Seagrove formation intermediate to mafic volcanic and volcanoclastic rocks (Cambrian to Neoproterozoic)—Map scale bodies of metamorphosed andesitic to basaltic lavas and shallow intrusions (like CZsimv) with interlayered volcanoclastic rocks consisting of phyllitic siltstone to conglomerate.
CZsmd	Seagrove formation mudstones (Cambrian to Neoproterozoic)—Grayish-green, light gray to dark gray, locally with distinctive reddish-gray or maroon to lavender coloration, metamorphosed, thinly bedded

mudstone, siltstone and sandy siltstone. Mudstone bedding ranges from mm-scale up to 10 cm, bedding layers traceable for several feet locally, may exhibit soft sediment deformation. Locally tuffaceous with a relict vitric texture. When present, crystal fragments are usually plagioclase with rare quartz. Deposition interpreted as distal from volcanic center.

CZse/pl

Seagrove formation epiclastic and pyroclastic rocks with rhyodacitic

domes of the Robbins area (Cambrian to Neoproterozoic)—Gray to greenish-gray, metamorphosed tuffaceous sandstones, conglomeratic sandstones, siltstones and local interbeds of mudstone. Contains lesser amounts of fine- to coarse tuff, welded tuff and rhyodacitic lavas/shallow intrusions. The siltstones typically are weakly phyllitic. Fiamme-like shaped clasts are common in the conglomeratic sandstones, sandstones and tuffs. Quartz and feldspar crystal fragments are common in the sedimentary components, tuffs and lavas. Silicified and/or sericitized altered rock and quartz veining present. Interpreted as a felsic sedimentary package deposited contemporaneous with local felsic volcanism. Correlates to Unit 2 of Powers (1993).

CZshar

Seagrove formation hydrothermally altered rocks (Cambrian to

Neoproterozoic)—Areas within the Seagrove formation of moderate to intense hydrothermally altered rock. Alteration consists of silicified, sericitized and pyrophyllitized rock. Chloritoid locally present. Sulfides are common. Primary sedimentary or volcanic textures are commonly

obliterated. Includes the moderate argillic (sericite) alteration zone in the Robbins area as described by Powers (1993).

CZshar-pqs **Pyrophyllite-quartz-sericite hydrothermally altered rocks (Cambrian to Neoproterozoic)**—Zones of intense hydrothermally altered rock with pyrophyllite. Include zones of pyrophyllite-quartz +-sericite alteration. Primary sedimentary or volcanic textures are obliterated. Includes active, inactive and abandoned pyrophyllite mines and prospects. After Powers (1993).

CZshar-qsh **Quartz-sericite-hematite hydrothermally altered rocks (Cambrian to Neoproterozoic)**—Zones of intense hydrothermally quartz-sericite-hematite altered rock. Includes silicified rock that may form resistant ridges in the Robbins area. According to Powers (1993), hematitic rocks have low to no detection of gold.

CZshar-qsp **Quartz-sericite-pyrite hydrothermally altered rocks (Cambrian to Neoproterozoic)**—Zones of intense hydrothermally quartz-sericite-pyrite altered rock. Includes silicified rock that may form resistant ridges in the Robbins area. According to Powers (1993), pyritic rocks have moderate (>100 ppb) to high (>1000 ppb) gold occurrence.

UWHARRIE VOLCANIC COMPLEX

CZurlt **Uwharrie Volcanic Complex rhyodacitic lavas and tuffs (Cambrian to Neoproterozoic)**—Gray to dark gray, siliceous, aphanitic rhyodacites and tuffs, quartz-bearing tuffs and quartz-rhyodacite porphyry. Rhyodacites and tuffs are vitric-like and weakly to moderately foliated. Rhyodacites are locally flow layered and(or) spherulitic. Phenocrysts of

albite are generally present, commonly accompanied by quartz. The size and abundance of phenocrysts varies widely. Tuffs locally contain fiamme-shaped clasts. May include lava flows, domes, necks , and dikes. Zuf, Felsite unit, of Seiders (1981).

CZufc **Felsic volcaniclastic rocks (Cambrian to Neoproterozoic)**—Thin- to very thick bedded, light- to dark-gray, in part pale-green, tuff, lapilli tuff, and tuff breccia. The most abundant rock types are very thick bedded tuff, lapilli tuff, and tuff breccia, but in many parts of the map unit subordinate amounts of well-bedded, thin- to thick-bedded, in part graded, tuff, tuffaceous sandstones, tuffaceous conglomeratic sandstones and felsic volcanic mudstone are interbedded. In many rocks the principal clasts are dense felsite, like that occurring in the numerous felsite bodies. Clasts are angular to subrounded, very rarely well rounded. Other rocks have clasts with poorly preserved textures that are strongly suggestive of pumice. These rocks generally lack interbeds of well-bedded tuff and may represent thick ashflow deposits. The probable pumice fragments are commonly much flattened but the flattening is in the plane of the regional cleavage or foliation and may be tectonic. Textural details are too poorly preserved to record most features that would clearly indicate welding.

CZuimcl **Intermediate to mafic volcaniclastic rocks and lavas (Cambrian to Neoproterozoic)**—Thin- to very thick-bedded, grayish-green tuff, lapilli tuff, and tuff breccia. Local occurrence of andesitic to basaltic lavas with hyaloclastic textures. Commonly contains scattered clasts of

felsic volcanic rock. Modified equivalent of CZumc of Seiders (1981).

CZuiml **Intermediate to mafic lavas (Cambrian to Neoproterozoic)**—Small bodies of metamorphosed andesite to basalt exposed chiefly in the northeastern part of the Seiders (1981) map area. Commonly amygdaloidal. Pillow lava occurs northeast of Asheboro, along the dead end road 1.5 km east of Oakie Mountain. May include some shallow intrusive rocks. Modified equivalent of CZuml of Seiders (1981).

CZuhar **Uwharrie Volcanic Complex hydrothermally altered rocks (Cambrian to Neoproterozoic)**—Areas within the Uwharrie Formation of moderate to intense hydrothermally altered rock. Alteration consists of silicified, sericitized and pyrophyllitized rock. Chloritoid locally present. Sulfides are common. Primary sedimentary or volcanic textures are usually obliterated.

ELLERBE AREA VOLCANICS AND VOLCANICLASTIC ROCKS

CZemd **Metamorphosed thin- to thickly-bedded mudstones of the Ellerbe area (Cambrian to Neoproterozoic)**—Green, grayish-green to gray, thin- to thickly-bedded (typically >1 cm thick) metamorphosed mudstone and siltstone ranging from felsic to mafic source material. Local occurrence of laminated mudstone and siltstone. Local interlayers of felsic tuff or tuffaceous siltstone present. Maybe related to the Cid Formation. New and unpublished LA-ICP-MS U-Pb detrital zircon ages from the unit include: maximum depositional ages (MDA) of 540 Ma for sample SP100K 551983 and MDA of 562 Ma for sample SP100K 551641.

CZeimvc **Metamorphosed intermediate to mafic volcanics and clastic rocks of**

the Ellerbe area (Cambrian to Neoproterozoic)—Metamorphosed intermediate to mafic volcanoclastic rocks and lavas with lesser mudstone and siltstone. Andesites and basalts locally display hyaloclastic and vuggy (interpreted as vesicles) texture. Local felsic sandstones present (similar in appearance to sandstones of the Seagrove formation). Likely a part of CZemd unit but with abundant intermediate to mafic volcanics and coarser-grained sediments.

CZelmd **Metamorphosed laminated mudstones of the Ellerbe area (Cambrian to Neoproterozoic)**—Grayish-green to gray, laminated to thin bedded (typically 1 mm to 1 cm thick) metamorphosed mudstone and siltstone. Beds are very regular and can be traced across the outcrop. Local interlayers of felsic tuff or tuffaceous siltstone present. Very similar in appearance to the Tillery Formation and may be equivalent. A new and unpublished LA-ICP-MS U-Pb detrital zircon age from the unit yields a maximum depositional age (MDA) of 543 Ma for sample SP100K 551786.

CONTACT AUREOLE OF THE LILESVILLE GRANITE

CZah-ex **Exterior aureole hornfels of the Lilesville Granite (Cambrian to Neoproterozoic)**—Mineralogy and color of rock varies with proximity to Lilesville Granite. In the extreme outer portions of the exterior aureole, the rocks are typical Carolina terrane metasediments with relict bedding with the presence of porphyroblasts of cordierite. As metamorphic grade increases closer to the granite, the rocks are hornfels that are grey, well-indurated with dark spots up to several millimeters (i.e. cordierite and other porphyroblasts). In the inner exterior aureole the

hornfels are black, dense and extremely well-indurated (may be confused with diabase). The mineral assemblage varies with location and may include cordierite, garnet, biotite, and other minerals. Relict bedding is locally preserved as faint striations on weathered surfaces. See Evans and Speer (1984) for detailed descriptions.

CZah-in **Interior aureole hornfels of the Lilesville Granite (Cambrian to Neoproterozoic)**—Gray to black, dense hornfels that vary in texture and mineral assemblages. Includes porphyroblasts of cordierite, garnet, biotite, scarce orthopyroxene, and other minerals. Locations with abundant biotite have a gneissic texture. See Evans and Speer (1984) for detailed descriptions.

CZah-mig **Migmatized interior aureole hornfels of the Lilesville Granite (Cambrian to Neoproterozoic)**—Gray to black, migmatized hornfels. Resemble polydeformed gneisses associated with regional metamorphism. Include zones of leucosomes, ptigmatically folded leucosome veins and boudins. Minerals include cordierite, garnet, biotite, sillimanite, and other minerals. See Evans and Speer (1984) for detailed descriptions.

HYCO ARC

HYCO FORMATION - UPPER PORTION

HYDROTHERMALLY ALTERED UNITS

Zhhar **Hydrothermally altered rocks (Neoproterozoic)**—Mixed unit of hydrothermally altered rocks consisting of: dense siliceous

cryptocrystalline rock; quartz-pyrophyllite rocks, +- kaolinite, andalusite, chloritoid, sericite, paragonite and iron oxides; quartz-sericite rocks, +- paragonite, k-feldspar and iron oxides; and quartz-chloritoid-chlorite rocks, +- sericite and hematite. Described in detail by Hughes (1987) and Schmidt et al. (2006).

- Zhat/vcs **Altered tuffs and volcanoclastic sedimentary rocks (Neoproterozoic)—**
Mixed unit of altered volcanoclastic rocks and volcanoclastic sedimentary rocks. Alteration consists of silicified, sericitized and pyrophyllitized rock. Chloritoid locally present. Volcanoclastic sedimentary rocks include conglomeratic siltstone to conglomerate that may be variably altered. Includes area of quartz-sericite-paragonite rock (Zvqs) of Schmidt et al. (2006). Andesitic to basaltic lavas and massive quartz locally present.
- Zhe/pl-a **Altered mixed epiclastic-pyroclastic rocks with interlayered dacitic lavas (Neoproterozoic)—**Area of unit Zhe/pl that is mildly to moderately hydrothermally altered. Alteration consists of silicified, sericitized and pyrophyllitized rock. Intensity of alteration varies locally.

VOLCANOCLASTIC-SEDIMENTARY UNITS

- Zhe/pl **Mixed epiclastic-pyroclastic rocks with interlayered dacitic lavas (Neoproterozoic)—**Mixed epiclastic-pyroclastic rocks with interlayered dacitic lavas: Grayish-green to greenish-gray, locally with distinctive reddish-gray or maroon to lavender coloration; metamorphosed: conglomerate, conglomeratic sandstone, sandstone, siltstone and mudstone. Lithologies are locally bedded; locally tuffaceous with a

cryptocrystalline-like groundmass. Siltstones are locally phyllitic. Locally contain interbedded dacitic lavas identical to Zhdlt unit. Contains lesser amounts of fine- to coarse tuff and lapilli tuff with a cryptocrystalline-like groundmass. Pyroclastics, lavas, and epiclastics are mainly felsic in composition. Minor andesitic to basaltic lavas and tuffs present. Silicified and/or sericitized altered rock are locally present. Conglomerates and conglomeratic sandstones typically contain subrounded to angular clasts of dacite in a clastic matrix. Portions of the Zhe/pl unit are interpreted to have been deposited proximal to active volcanic centers represented by the Zhdlt unit but are also interpreted to record the erosion of proximal volcanic centers after cessation of active volcanism.

FELSIC TO INTERMEDIATE VOLCANIC UNITS

Zhdlt (u)

Dacitic lavas and tuffs of the upper portion of the Hyco Formation

(Neoproterozoic)—Greenish-gray to dark gray, siliceous, aphanitic dacite, porphyritic dacite with plagioclase phenocrysts, and flow banded dacite. Dacite with hyaloclastic textures are common. Welded and non-welded tuffs associated with the lavas include: greenish-gray to grayish-green, fine tuff, coarse plagioclase crystal tuff and lapilli tuff. Locally, interlayers of immature conglomerate and conglomeratic sandstone with abundant dacite clasts are present. The dacites are interpreted to have been coherent extrusives or very shallow intrusions associated with dome formation. The tuffs are interpreted as episodic pyroclastic flow deposits, air fall tuffs or reworked tuffs generated

during formation of dacite domes. The unit occurs as map scale pods surrounded by clastic rocks of Zhe/pl unit. Wortman et al. (2000) reports an age of $615.7 \pm 3.7/-1.9$ Ma U-Pb zircon date for a dacitic tuff from the unit in the Rougemont quadrangle in the Henderson 100K.

INTERMEDIATE TO MAFIC VOLCANOCLASTIC-SEDIMENTARY UNITS

Zhime/pl

Mixed intermediate to mafic epiclastic-pyroclastic rocks with

interlayered intermediate to mafic lavas (Neoproterozoic)—Grayish-green to green, locally with distinctive reddish-gray or maroon to lavender coloration; metamorphosed: conglomerate, conglomeratic sandstone, sandstone, siltstone and mudstone. Lithologies are locally bedded; locally tuffaceous with a cryptocrystalline-like groundmass. Siltstones are locally phyllitic. Locally contain interbedded intermediate to mafic lavas identical to Zhablt, Zhabl, and Zhablc units. Contains lesser amounts of fine- to coarse tuff and lapilli tuff with a cryptocrystalline-like groundmass. Pyroclastics, lavas, and epiclastics are mainly intermediate to mafic in composition. Minor dacitic lavas and tuffs present. Silicified and/or sericitized altered rock similar to Zhat unit are locally present. Conglomerates and conglomeratic sandstones typically contain subrounded to angular clasts of andesite and basalt in a clastic matrix. Generally interpreted to have been deposited proximal to active intermediate to mafic composition volcanic centers and/or record the erosion of proximal intermediate to mafic composition volcanic centers after cessation of active volcanism. May be related to Green et al. (1982) unit C - Intermediate to Mafic

Volcanics and Graywacke.

INTERMEDIATE TO MAFIC VOLCANIC UNITS

Zhabl **Andesitic to basaltic lavas (Neoproterozoic)**—Green, gray-green, gray, dark gray and black; typically unfoliated, amygdaloidal, plagioclase porphyritic, amphibole/pyroxene porphyritic and aphanitic; andesitic to basaltic lavas and shallow intrusions. Hyaloclastic texture is common and imparts a fragmental texture on some outcrops and float boulders. Conglomeratic rocks consisting of angular clasts of andesite and/or basalt occur locally and are interpreted as resedimented hyaloclastite.

Zhable **Andesitic to basaltic lavas with interlayered epiclastic rocks (Neoproterozoic)**—Light green, gray-green, gray, and dark gray; typically unfoliated, amygdaloidal, plagioclase porphyritic, amphibole/pyroxene porphyritic and aphanitic; metamorphosed: andesitic to basaltic lavas and shallow intrusions. Hyaloclastic texture is common and imparts a fragmental texture on some outcrops and float boulders. Contains lesser amounts of grayish-green, light green, and light gray to white; metamorphosed conglomerate, conglomeratic sandstone, sandstone, siltstone and mudstone.

REFERENCES:

- Barefoot, J., 2015, Chronological and Mineralogical Comparison of Gold Prospect Host Rock with Known Gold Horizon Host Rock - Orange and Chatham Counties, North Carolina, unpublished senior thesis, University of North Carolina at Chapel Hill, 18 pages.
- Bartlett, C.S., Jr. 1967, Geology of the Southern Pines Quadrangle [MS Thesis], University of North Carolina at Chapel Hill, 101 p.

Black, W.W., 1986, Simple Bouguer gravity map of North Carolina and vicinity: North Carolina Geological Survey Open-File Report 87-1, map scale 1:500,000, 1 sheet, descriptive text, 3 p.

Blake, D.E., Clark, T.W., and Heller, M.J., 2001, A Temporal View of Terranes and Structures in the Eastern North Carolina Piedmont, *in* Field Trip Guide, Southeastern Section - Geological Society of America, 2001, p. 149-177.

Blake, D.E., Stoddard, E.F., Bradley, P.J., and Clark, T.W., 2012, Neoproterozoic to Mesozoic petrologic and ductile-brittle structural relationships along the Alleghanian Nutbush Creek fault zone and Deep River Triassic basin in North Carolina, *in* Eppes, M.C., and Bartholomew, M.J., eds., From the Blue Ridge to the Coastal Plain: Field Excursions in the Southeastern United States: Geological Society of America Field Guide 29, p. 219–261.

Blake, D.E., Stoddard, E.F., Bradley, P.J., and Pelt, K.E. 2025, Compiled bedrock geologic map of the Henderson and western portion of the Roanoke Rapids 30' x 60' quadrangles, North Carolina: North Carolina Geological Survey, Open-file Report 2025-04, scale 1:100,000, in color, with accompanying pamphlet.

Bobyarchick, A.R., 1981, The Eastern Piedmont fault system and its relationship to Alleghanian tectonics in the southern Appalachians: *The Journal of Geology*, v. 89, p. 335–347, doi:10.1086/628595.

Boorman, S. L., Brennan, M.P., Oliver, J.A., Hibbard, J.P., and Fodor, R. V., 2013, Field relationship between the Uwharrie Formation and the lower Albemarle Group and geochemistry of the 'Morrow Mountain Rhyodacites', central North Carolina: implications for the stratigraphy of the Albemarle arc, *in* Hibbard, J.P., and Pollock, J.C., eds., One arc, two arcs, old arc, new arc: the Carolina terrane in central North Carolina, Carolina Geological Society field trip guidebook, p. 155-173.

Bowman, J.D., 2010, The Aaron Formation: Evidence for a New Lithotectonic Unit in Carolina, North Central North Carolina, unpublished M.S. thesis, North Carolina State University, Raleigh, North Carolina, 116 p.

Bowman, J.D., Hibbard, J.P. and Miller, B.V. 2013, The Virgilina Sequence Redefined, North Central North Carolina, *in* Hibbard, J.P., and Pollock, J.C., eds., One arc, two arcs, old arc, new arc: the Carolina terrane in central North Carolina, Carolina Geological Society field trip guidebook, p. 127-138.

Bradley, P.J., and Miller, B.V., 2011, New geologic mapping and age constraints in the Hyco Arc of the Carolina terrane in Orange County, North Carolina: Geological Society of America Abstracts with Programs, Vol. 43, No. 2.

Bradley, P.J. and Bolich, R.E., 2022, Groundwater features map of Chatham County and surrounding areas, North Carolina: North Carolina Geological Survey Open-file Report 2022-04, scale 1:50,000, in color. <https://deq.nc.gov/energy-mineral-and-land-resources/geological-survey/ofrs-geological-survey/groundwater-features-map-chatham-county-and-surrounding-areas-north-carolina>

Bradley, P.J., 2023, Geology of Chatham County and Surrounding Areas, North Carolina, *in* Bradley, P.J., ed., The geology of the Carolina terrane in Chatham County, North Carolina, Carolina Geological Society annual meeting field trip guidebook for 2023, p. 1-26.

Bradley, P.J., Thompson, G.J. and Pelt, K.E., 2025a, Compiled Bedrock Geologic Map of the Chapel Hill 30' x 60' Quadrangle (Chapel Hill 100K), North Carolina, North Carolina Geological Survey Open-file Report 2025-06, scale 1:100,000, in color, with accompanying pamphlet.

Bradley, P.J., Curry, A.C., Thompson, G.J. and Zhang, Q., 2025b, New problems from new data - an example from the Carolina terrane in North Carolina and bearing on interstate correlation, Geological Society of America Abstracts with Programs., Vol. 57, No. 2, 2025, doi: 10.1130/abs/2025SE-407549

Brazell, S.J., 2013, Sedimentology and depositional environments of the Wadesboro sub-basin, eastern Piedmont, North Carolina, [M.S. Thesis], University of North Carolina Charlotte, 196 p.

Butler, J.R., and Secor, D.T., Jr., 1991, The central Piedmont, *in* Horton, W., and Zullo, V., eds., The Geology of the Carolinas: Knoxville, University of Tennessee Press, p. 59–78.

Campbell, M.R., and Kimball, K.W., 1923, The Deep River coal field of North Carolina: North Carolina Geological and Economic Survey Bulletin 33, 95 p.

Campbell, S.K., 1984, Geology, petrography, and geochemistry of the Lemon Springs pluton and associated rocks, Lee County, North Carolina [M.S. Thesis]: East Carolina University, Greenville, North Carolina, 132 p.

Carpenter, P.A., 1999, Bedrock Geologic map of the Erect 7.5-minute Quadrangle, Randolph County, North Carolina, North Carolina Geological Survey, NCGS Open-file Report 1999-01, scale 1:24,000.

Clark, T.W., Gore, P.J., and Watson, M.E., 2001, Depositional and structural framework of the Deep River Triassic basin, North Carolina, *in* Hoffman, C.W., ed. Field Trip Guidebook for the 50th Annual Meeting of the Southeastern Section, Geological Society of America, Raleigh, North Carolina, p. 27-50. (re-printed in Carolina Geological Society Field Trip Guidebook 2011)

Clark, T.W., Taylor, K.B., Bradley, P.J., 2011, Geology, natural gas potential and mineral resources of Lee, Chatham and Moore counties, North Carolina, Carolina Geological Society field trip guidebook for 2011, 49 p.

Conley, J.F., 1962a, Geology of the Albemarle Quadrangle, North Carolina, North Carolina, Division of Mineral Resources, North Carolina Department of Conservation and Development (North Carolina Geological Survey), Bulletin 75, 26 p.

Conley, J.F., 1962b, Geology and mineral resources of Moore County, North Carolina: Division of Mineral Resources, North Carolina Department of Conservation and Development, Bulletin 76, 40 p.

Conley, J.F., and Bain, G.F., 1965, Geology of the Carolina Slate belt west of the Deep River-Wadesboro Triassic basin, North Carolina, *Southeastern Geology* v. 6, no. 3, p. 117-138

Daniels, D.L., and Leo, G.W., 1985, Geologic interpretation of basement rocks of the Atlantic Coastal Plain: U.S. Geological Survey Open-File Report 85-655, 45 p.

DeDecker, J., Coleman, D.S., Hibbard, J.P., and Pollock, J.C., 2013, Preliminary U-Pb TIMS zircon ages from the Stony Mountain gabbro at Ridges Mountain, North Carolina: timing of the birth of the Rheic Ocean? *In*, Hibbard, J.P. and Pollock, J., (Eds.). One arc, two arcs, old arc, new arc: The Carolina terrane in central North Carolina. Carolina Geological Society Guidebook, p. 239-244.

- Dineen, M.J., 1982, Geology of the Triassic "Ellerbe Basin", Richmond County, North Carolina, unpublished M.S. thesis, North Carolina State University, Raleigh, 68 p.
- Evans, N. H. and Speer, J. A., 1984, Low-pressure metamorphism and anatexis of Carolina Slate belt phyllites in the contact aureole of the Lilesville pluton, North Carolina, USA, *Contributions to Mineralogy and Petrology* v. 87, p. 297-309.
- Farrar, S.S., 1985, Stratigraphy of the northeastern North Carolina Piedmont: *Southeastern Geology*, v. 25, no. 3, p. 159-183.
- Fullagar, P. D. and Butler, J. R., 1979, 325 to 265 m.y. old granitic plutons in the Piedmont of the southeastern Appalachians: *American Journal of Science*, v. 279, p. 161-185.
- Goldberg, S. A., 1994, U-Pb geochronology of volcanogenic terranes of the eastern North Carolina Piedmont: preliminary results, *in* Stoddard, E.F. and Blake, D. E., eds., *Geology and Field Trip Guide, Western Flank of the Raleigh metamorphic belt, North Carolina*, Raleigh, North Carolina Geological Survey, Carolina Geological Society Guidebook for 1994, p. 13-17.
- Goldsmith, Richard, Milton, D.J., and Horton, J.W., 1988, Geologic map of the Charlotte 1 degree x 2 degree quadrangle, North and South Carolina: U.S. Geological Survey, Miscellaneous Investigations Series Map I-1251-E, scale 1:250000.
- Glover, L., and Sinha, A., 1973, The Virgilina deformation, a late Precambrian to Early Cambrian (?) orogenic event in the central Piedmont of Virginia and North Carolina, *American Journal of Science*, Cooper v. 273-A, pp. 234-251.
- Green, G., Cavaroc, V., Stoddard, E., Abdelzahir, A., 1982, Volcanic and volcanoclastic facies in a part of the slate belt of North Carolina, In: Bearce, D., Black, W., Kish, S., Tull, J. (Eds.), *Tectonic studies in the Talladega and Carolina slate belts, Southern Appalachian Orogen*. Geological Society of America Special Paper, vol. 191, pp.109– 124.
- Grimley, D.A., 2023, Quaternary terraces and deposits in Chatham County area, North Carolina, *in* Bradley, P.J., ed., *The geology of the Carolina terrane in Chatham County, North Carolina*, Carolina Geological Society annual meeting field trip guidebook for 2023.
- Harris, C.W., 1984, Coarse-grained submarine-fan deposits of magmatic arc affinity in the late Precambrian Aaron Formation, North Carolina, U.S.A., *Precambrian Research*, 26, pp. 285-306.
- Harris, C., and Glover, L., 1985, The Virgilina deformation: implications of stratigraphic correlation in the Carolina slate belt, *Carolina Geological Society field trip guidebook*, 36 p.
- Harris, C., and Glover, 1988, The regional extent of the ca. 600 Ma Virgilina deformation: implications of stratigraphic correlation in the Carolina terrane, *Geological Society of America Bulletin*, v. 100, pp. 200-217.
- Hatcher, R.D., Howell, D., and Talwani, P., 1977, Eastern Piedmont fault system: Speculation on its extent: *Geology*, v. 5, p. 636–640, doi:10.1130/0091-7613(1977)5<636:EPFSSO>2.0.CO;2.

Hibbard, J., and Samson, S., 1995, Orogenesis exotic to the Iapetan cycle in the southern Appalachians, In, Hibbard, J., van Staal, C., Cawood, P. editors, Current Perspectives in the Appalachian– Caledonian Orogen. Geological Association of Canada Special Paper, v. 41, pp. 191–205.

Hibbard, J., Stoddard, E.F., Secor, D., Jr., and Dennis, A., 2002, The Carolina Zone: Overview of Neoproterozoic to early Paleozoic peri-Gondwanan terranes along the eastern flank of the southern Appalachians: *Earth Science Reviews*, v. 57, n. 3/4, p. 299-339.

Hibbard, J. P., van Staal, C. R., Rankin, D. W., and Williams, H., 2006, Lithotectonic map of the Appalachian Orogen, Canada-United States of America, Geological Survey of Canada, Map-2096A. 1:1,500,000-scale.

Hibbard, J.P., van Staal, C.R., and Rankin, D.W., 2010, Comparative analysis of the geological evolution of the northern and southern Appalachian orogen: Late Ordovician-Permian: in Tollo, R.P., Bartholomew, M.J., Hibbard, J.P., and Karabinos, P.M., eds., From Rodinia to Pangea: The Lithotectonic Record of the Appalachian Region: Geological Society of America Memoir 206, p. 51-69.

Hibbard, J.P., Miller, B.V., Allen, J.S., Standard, I.D., Hames, W.E., Lavalley, S.B., and Boland, I.B., 2012, Kinematics, U-Pb geochronology, and ⁴⁰Ar/³⁹Ar thermochronology of the Gold Hill shear zone, North Carolina: the Cherokee orogeny in Carolinia, southern Appalachians. *Geological Society of America Bulletin*, v. 124, p.643-656.

Hibbard, J.P., Pollock, J.C., and Bradley, P.J., 2013, One arc, two arcs, old arc, new arc: An overview of the Carolina terrane in central North Carolina, Carolina Geological Society field trip guidebook, 265 p.

Hughes, E.H., 1987, The geology and hydrothermal alteration centers of the Snow Camp Mine-Major Hill area, central Carolina slate belt, Alamance and Chatham Counties, North Carolina: U.S. Geological Survey, Open-File Report OF-87-180, scale 1:24000.

Ingle, S., Mueller, P., Heatherington, A., and Kozuch, M., 2003, Isotopic evidence for the magmatic and tectonic histories of the Carolina terrane: implications for stratigraphy and terrane affiliation, *Tectonophysics*, 371, p. 187-211.

Kish, S.A., and Campbell, S.K., 1986, A middle Paleozoic plutonic terrane in the eastern Piedmont of North Carolina: *Geological Society of America Abstracts with Programs*, v. 18, no. 6, pg. 658.

Klein, T.L., 1985, Glendon Pyrophyllite deposits – Stops 2-10, in Feiss, P.G., editor, Volcanic-hosted gold and high-alumina rocks of the Carolina slate belt, guidebook for the field trip held in conjunction with the 1985 fall meeting of the Society of Economic Geologists and the 1985 annual meeting of the Geological Society of America, Orlando, Florida, p. 48-72.

Lawrence, D., and Hoffman, C.H., 1993, Geology of basement rocks beneath the North Carolina Coastal Plain. North Carolina Geological Survey, Bulletin 95, 60 p.

Milton, D., 1984, Revision of the Albemarle Group, North Carolina: U.S. Geological Survey Bulletin 1537-A, p. A69–A72.

Moore, J. G. and Lockwood, J. P., 1973, Origin of comb layering and orbicular structure, Sierra Nevada batholith, California: *Geological Society of America Bulletin*, v. 84, p. 1-19.

Morrison, D.J., and Coleman, D, 2023, Absolute age determination of the Parks Crossroads granodiorite of the Carolina terrane, *in* Bradley, P.J., ed., The geology of the Carolina terrane in Chatham County, North Carolina, Carolina Geological Society annual meeting field trip guidebook for 2023.

Moye, R.J., 2023a, Introduction to: The Albemarle Sequence of the Carolina Terrane in Central North Carolina: Geologic and metallogenic Analysis with an alternative model, *in* Bradley, P.J., ed., The geology of the Carolina terrane in Chatham County, North Carolina, Carolina Geological Society annual meeting field trip guidebook for 2023.

Moye, R.J., 2023b, The Albemarle Sequence of the Carolina Terrane in Central North Carolina: Geologic and metallogenic analysis with an alternative model, *in* Supplemental Papers, Carolina Geological Society Annual Meeting 2023, p. 1-82.

Moye, Robert J., 2018, Star and Carter gold mines, Montgomery County, North Carolina: Review of geology and metallogeny: North Carolina Geological Survey, Open-File Report 2018-17, 33p.

NCDOT, 2026, North Carolina Department of Transportation Geotechnical Engineering Unit, Boring Group Webmap, ArcGIS Map Service, <https://ncdot.maps.arcgis.com/apps/mapviewer/index.html?webmap=00d09c8cf5034b5a9227ff5aa09f3e14> (Accessed in Spring 2026).

Olsen, P. E., Froelich, A J., Daniels, D. L., Smoot, J. P., and Gore, P. J. W., 1991, Rift basins of early Mesozoic age, *in* Horton, W., ed., Geology of the Carolinas, University of Tennessee Press, Knoxville, p. 142-170.

Parker, J.M., 1979, Geology and mineral resources of Wake County: North Carolina Geological Survey Bulletin 86, 122 p., 1:100,000-scale map.

Pelt, K.E. and Bradley, P.J., 2023, Preliminary detrital zircon data from Chatham County, NC, *in* Bradley, P.J., ed., The geology of the Carolina terrane in Chatham County, North Carolina, Carolina Geological Society annual meeting field trip guidebook for 2023.

Pollock, J. C., 2007, The Neoproterozoic-Early Paleozoic tectonic evolution of the peri-Gondwanan margin of the Appalachian orogen: an integrated geochronological, geochemical and isotopic study from North Carolina and Newfoundland. Unpublished PhD dissertation, North Carolina State University, 194 p.

Pollock, J.C., Hibbard, J.P., and Sylvester, P.J., 2010, Depositional and tectonic setting of the Neoproterozoic-early Paleozoic rocks of the Virgilina sequence and Albemarle Group, North Carolina: *in* Tollo, R.P., Bartholomew, M.J., Hibbard, J.P., and Karabinos, P.M., eds., From Rodinia to Pangea: The Lithotectonic Record of the Appalachian Region: Geological Society of America Memoir 206, p. 739-772.

Powers, J.A., 1993, Gold Mineralization and Associated Hydrothermal Alteration in the Robbins District, Moore County, North Carolina: Unpublished PhD dissertation, University of North Carolina, Chapel Hill, North Carolina, 172 p.

Randazzo, A.F., and Copeland, R.E., 1976, The geology of the northern portion of the Wadesboro Triassic basin, North Carolina, *Southeastern Geology*, v. 17, p. 115-138.

- Randazzo, A., 1972, Petrography and stratigraphy of the Carolina Slate Belt, Union County, North Carolina: North Carolina Department of Natural and Economic Resources, Raleigh, North Carolina, Special Publication No. 4, 38p.
- Reading, H. G., and Richards, M., 1994, Turbidite systems in deep-water Basin margins classified by grain size and feeder system, *Bulletin American Association Petroleum Geologists*, 78, 792-822.
- Reinemund, J.A., 1955, Geology of the Deep River coal field, North Carolina: U.S. Geol. Survey Prof. Paper 246, 159 p.
- Schmidt, R., 1985, High-alumina hydrothermal systems in volcanic rocks and their significance to mineral prospecting in the Carolina slate belt. U.S. Geological Survey Bulletin 1562, 59 p.
- Schmidt, R.G., Gumiel, P., and Payas, A., 2006, Geology and mineral deposits of the Snow Camp-Saxapahaw area, Central North Carolina: United States Geological Survey Open-file Report 2006-1259 (<http://pubs.usgs.gov/of/2006/1259/index.html>).
- Seiders, V. M., and Wright, J. E., 1977, Geology of the Carolina volcanic slate belt in the Asheboro, North Carolina, area, p. 1-34, *in* Burt, E. R., ed., Field guides for Geological Society of America, Southeastern Section Meeting, Winston-Salem, North Carolina Department of Natural and Economic Resources, Geology and Mineral Resources Section, 149 p.
- Seiders, V., 1981, Geologic map of the Asheboro Quadrangle, North Carolina, and adjacent areas. US Geological Survey, Miscellaneous Investigations Map I-1314, scale 1:62,500.
- Stoddard, E., Farrar, S., Horton, W., Butler, R., and Druhan, R., 1991, The eastern Piedmont in North Carolina, *in* Horton, J. W., Jr., and Zullo, V. A., eds., *The Geology of the Carolinas*: Knoxville, Tennessee, University of Tennessee Press, p. 79–92.
- Stow, D.A.V. and Mayall, M., editors, 2000, Deep-water Sedimentary Systems: Thematic Set, Marine and Petroleum Geology, Volume 17, No. 2.
- Stromquist, A., and Henderson, J., 1985, Geologic and geophysical maps of south-central North Carolina: U.S. Geological Survey Miscellaneous Investigation Series Map I-1400.
- Stromquist, A.A., and Sundelius, H.W., 1969. Stratigraphy of the Albemarle Group of the Carolina slate belt in central North Carolina: U.S. Geological Survey Bulletin 1274-B, p. 1-22.
- Stuckey, J.L., 1928, Pyrophyllite deposits of North Carolina: North Carolina Department of Conservation and Development, Bulletin 37, 62 p.
- USGS Piedmont Bridge, 2025, Piedmont Bridge South Magnetic and Radiometric Survey, Preliminary data provided to the NCGS.
- Upchurch, C.N., 1968, The Geology of the Southwest Quarter of the Troy Quadrangle, North Carolina, unpublished M.S. thesis, North Carolina State University, Raleigh, North Carolina 90 p.
- Waskom, John D.; Butler, J. Robert, 1971, Geology and Gravity of the Lilesville Granite Batholith, North Carolina, *GSA Bulletin*. 82 (10): 2827–2844.
- Waskom, J. D., 1970, Geology and Geophysics of the Lilesville Granite Batholith, North

Carolina: Ph.D. dissertation, University of North Carolina at Chapel Hill, 78 p.

Wortman, G.L., Samson, S.D., and Hibbard, J.P., 2000, Precise U-Pb zircon constraints on the earliest magmatic history of the Carolina terrane, *Journal of Geology*, v. 108, pp. 321-338.

Zietz, I., Riggle, F.E., and Gilbert, F.P., 1984, Aeromagnetic map of North Carolina: U.S. Geological Survey Geophysical Investigations Map GP-957, 1 sheet, scale 1:500,000.

Appendix A

Geochronology plots and tables

Geochronology data reduction and interpretation by Adam C. Curry


Zircon U-Pb isotope data were reduced using IsoplotR to assess degree of concordance prior to calculating a weighted mean crystallization (or metamorphic) age for igneous/metamorphic samples or estimating a maximum depositional age (MDA) for metasedimentary ages (Vermeesch, 2018, 2021a, 2021b). We used the Concordia distance filter (d_c) of Vermeesch (2021a) to filter data for degree of concordance, discarding values outside of the range $-2 < d_c < 4.6$, in addition to visual assessment of concordance, discarding any analyses not touching Concordia. Zircon U-Pb analyses were also excluded in cases where Pb loss was apparent (too young ages) or Th/U ratios were anomalous. Igneous crystallization ages reported here are zircon $^{238}\text{U}/^{206}\text{Pb}$ weighted mean ages, with errors reported at a 95% confidence interval. Outliers were not included in the weighted mean ages, using Chauvenet's Criterion in IsoplotR. Estimations of MDAs for detrital zircon ages are based on the Maximum Likelihood Age (MLA) developed by Galbraith and Laslett (1993) for fission track thermochronology and extended to zircon U-Pb detrital zircon geochronology by Vermeesch (2021b).

Regarding Th/U ratios in zircon, a value of 0.1 has typically been the cutoff for igneous ($\text{Th}/\text{U} > 0.1$) and metamorphic ($\text{Th}/\text{U} < 0.1$) zircons (Rubatto, 2017). However, metamorphic zircons can range from <0.01 to >10 , whereas igneous zircons rarely have $\text{Th}/\text{U} < 0.1$ (Yakymchuk et al., 2018). Therefore, to isolate a metamorphic signature in a zircon population, it is necessary to look at zircons with $\text{Th}/\text{U} < 0.1$.

References


- Galbraith, R. F., & Laslett, G. M. (1993). Statistical models for mixed fission track ages. *Nuclear Tracks and Radiation Measurements*, 21(4), 459–470. [https://doi.org/10.1016/1359-0189\(93\)90185-C](https://doi.org/10.1016/1359-0189(93)90185-C)
- Gehrels, G. E., Valencia, V. A., & Ruiz, J. (2008). Enhanced precision, accuracy, efficiency, and spatial resolution of U-Pb ages by laser ablation-multicollector-inductively coupled plasma-mass spectrometry. *Geochemistry, Geophysics, Geosystems*, 9(3), 1–13. <https://doi.org/10.1029/2007GC001805>
- Gehrels, G., & Pecha, M. (2014). Detrital zircon U-Pb geochronology and Hf isotope geochemistry of Paleozoic and Triassic passive margin strata of western North America. *Geosphere*, 10(1), 49–65. <https://doi.org/10.1130/GES00889.1>
- Horstwood, M. S. A., Košler, J., Gehrels, G., Jackson, S. E., McLean, N. M., Paton, C., Pearson, N. J., Sircombe, K., Sylvester, P., Vermeesch, P., Bowring, J. F., Condon, D. J., & Schoene, B. (2016). Community-Derived Standards for LA-ICP-MS U-(Th)-Pb Geochronology – Uncertainty Propagation, Age Interpretation and Data Reporting. *Geostandards and Geoanalytical Research*, 40(3), 311–332. <https://doi.org/10.1111/j.1751-908X.2016.00379.x>
- Pullen, A., Ibáñez-Mejía, M., Gehrels, G. E., Giesler, D., & Pecha, M. (2018). Optimization of a Laser Ablation-Single Collector-Inductively Coupled Plasma-Mass Spectrometer (Thermo Element 2) for Accurate, Precise, and Efficient Zircon U-Th-Pb Geochronology. *Geochemistry, Geophysics, Geosystems*, 19(10), 3689–3705. <https://doi.org/10.1029/2018GC007889>
- Rubatto, D. (2017). Zircon: The Metamorphic Mineral. *Reviews in Mineralogy and Geochemistry*, 83, 261–295.
- Sundell, K. E., Gehrels, G. E., & Pecha, M. E. (2021). Rapid U-Pb Geochronology by Laser Ablation Multi-Collector ICP-MS. *Geostandards and Geoanalytical Research*, 45(1), 37–57. <https://doi.org/10.1111/ggr.12355>
- Vermeesch, P. (2018). IsoplotR: A free and open toolbox for geochronology. *Geoscience Frontiers*, 9(5), 1479–1493. <https://doi.org/10.1016/j.gsf.2018.04.001>
- Vermeesch, P. (2021a). Maximum depositional age estimation revisited. *Geoscience Frontiers*, 12(2), 843–850. <https://doi.org/10.1016/j.gsf.2020.08.008>
- Vermeesch, P. (2021b). On the treatment of discordant detrital zircon U-Pb data. *Geochronology*, 3(1), 247–257. <https://doi.org/10.5194/gchron-3-247-2021>
- Yakymchuk, C., Kirkland, C. L., & Clark, C. (2018). Th/U ratios in metamorphic zircon. *Journal of Metamorphic Geology*, 36(6), 715–737. <https://doi.org/10.1111/jmg.12307>

NCGS Station SP100K – 550027
Rhyodacite (metamorphosed)
Crystallization age



Sample SP100K-550027 has outlying ages of 506.5 ± 14.1 Ma, 560.6 ± 17.6 Ma, 563.2 ± 21.7 Ma, 571.5 ± 16.1 Ma, and 575.1 ± 25.4 Ma.

NCGS Station SP100K – 550264
Rhyodacite (metamorphosed)
Crystallization age




Sample SP100K-550264 has two outlying ages of 580.3 ± 29.4 Ma and 581.7 ± 15.9 Ma.

NCGS Station SP100K-550946

Tuffaceous pebbly sandstone (metamorphosed)

Detrital age




The MDA for sample SP100K-550946 is 523.8 ± 3.7 Ma ($n = 175$).


NCGS Station SP100K-550363

Conglomeratic sandstone (metamorphosed)


Detrital age



The MDA for sample SP100K-550363 is 542.8 ± 8.6 Ma ($n = 38$). It should be noted that this sample has a zircon age spectrum similar to an igneous crystallization age. It has low dispersion (see the ranked age plot for this sample). When a weighted mean age is calculated, it is 545.2 ± 2.5 Ma with an MSWD of 1.1, with two outliers ($n=36$).





NCGS Station SP100K-551063
Rhyodacite (metamorphosed)
Crystallization age



Sample SP100K-551063 (n=19) has a weighted mean age of 550.4 ± 4.2 Ma, with an MSWD of 0.44. Sample SP100K-551063 has an outlying age of 572 ± 15 Ma.

NCGS Station SP100K-551983
Laminated siltstone (metamorphosed)
Detrital age





Using all concordant analyses, the MDA for sample SP100K-551983 is 539.6 ± 2.2 Ma (n=241).

NCGS Station SP100K-551641


Siltstone (metamorphosed)

Detrital age




Using all concordant analyses older than 500 Ma, the MDA for sample SP100K-551641 is 561.5 ± 7.1 Ma. Multiple younger zircons were omitted from the MDA calculation. Using all concordant analyses, the MDA for sample SP100K-551641 is 339 ± 16 Ma ($n=47$). An MDA of ca. 339 Ma is interpreted as inaccurate and likely represents a metamorphic age. The sample location is proximal to the Lilesville granite - a ca. 330 Ma pluton.

NCGS Station SP100K-551786
 Sandy siltstone (metamorphosed)
 Detrital age



(n=205)



Using all concordant analyses older than 500 Ma, the MDA for sample SP100K-551786 is 542.7 ± 1.9 Ma. Multiple younger zircons were omitted from the MDA calculation. Using all concordant analyses, the MDA for sample SP100K-551786 is 326 ± 9.9 Ma (n=205). An MDA of ca. 326 Ma is interpreted as inaccurate and likely represents a metamorphic age. The sample location is proximal to the Lilesville granite - a ca. 330 Ma pluton.

Table 1. LA-ICP-MS U-Pb zircon data for the Southern Pines 100 K (crystallization ages)

Composition			Isotopic Ratios								Ages (Ma)						
ID	U (ppm)	U/Th	²⁰⁶ Pb/ ²⁰⁴ Pb	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (%)	²⁰⁷ Pb/ ²³⁵ U	Error (%)	²⁰⁶ Pb/ ²³⁸ U	Error (%)	error correlation	concordia distance (d _c) ^a	²⁰⁶ Pb/ ²³⁸ U	Error (Ma)	²⁰⁷ Pb/ ²³⁵ U	Error (Ma)	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (Ma)
Southern Pines 100k																	
550264																	
1 B	448	1.14	44894	17.2676	2.4	0.72023	3.2	0.090239	2.1	0.658	-1	557	11.1	550.8	13.5	525.5	52.3
2	454	1.13	60113	17.5307	2.8	0.70727	4.6	0.089966	3.7	0.791	-1.8	555.3	19.5	543.1	19.5	492.2	62.5
3 B	96	1.69	16792	17.3069	4.5	0.71247	6.6	0.08947	4.8	0.731	-0.99	552.4	25.5	546.2	27.9	520.4	98.8
5 A	3001	0.45	6871915	16.8844	1.2	0.73145	2.9	0.089611	2.7	0.907	0.71	553.2	14.1	557.4	12.6	574.4	27
16 D	111	2.44	10977	17.4929	4.4	0.70401	6.2	0.089358	4.3	0.698	-1.7	551.7	22.8	541.2	25.9	496.9	97.3
6 A	454	1.19	1018356	16.8792	2.2	0.72337	4.3	0.088595	3.8	0.867	0.8	547.2	19.7	552.7	18.5	575.1	47
6 B	520	1.32	212769	17.0452	2.7	0.70316	3.6	0.086966	2.4	0.667	0.54	537.6	12.4	540.7	15.1	553.8	58.6
6 E	532	0.9	62733	17.2087	1.9	0.72573	2.9	0.090619	2.2	0.765	-0.8	559.2	11.8	554	12.3	532.9	40.7
15 B	261	2.02	26412	17.4325	3.2	0.70996	4.4	0.089803	3	0.693	-1.6	554.4	16.2	544.7	18.5	504.6	69.7
15 C	449	2.01	26194	17.3234	2.4	0.70109	3	0.088125	1.8	0.592	-0.88	544.4	9.2	539.4	12.5	518.3	52.9
15 D	432	2.17	60104	16.8945	2.8	0.71776	3.7	0.087987	2.4	0.657	1	543.6	12.6	549.3	15.6	573.1	60.3
14 E	823	1.41	112202	17.2829	2.2	0.72355	3.2	0.090736	2.3	0.734	-1.1	559.9	12.5	552.8	13.6	523.5	47.4
14 F	812	1.15	19792	16.9523	2.3	0.74862	3.4	0.092084	2.5	0.734	-0.065	567.9	13.6	567.4	14.8	565.7	50.4
13 E	109	1.26	14537	17.3673	4.3	0.74746	6.8	0.094192	5.3	0.775	-2	580.3	29.4	566.7	29.7	512.8	94.8
7 A	181	2.24	8277	17.4627	3.1	0.71723	4.6	0.090879	3.4	0.735	-1.8	560.7	18.2	549	19.6	500.7	68.8
7 B	1308	1.01	152313	16.8846	1.8	0.72184	2.9	0.088436	2.3	0.777	0.83	546.3	11.9	551.8	12.5	574.4	40.2
8 D	209	2.88	57456	17.3162	3.3	0.71387	4.2	0.089695	2.5	0.607	-1.2	553.7	13.5	547	17.7	519.3	73.1
8 A	411	2.19	86621	16.936	1.9	0.71115	3.9	0.087434	3.4	0.87	0.79	540.3	17.7	545.6	16.6	567.8	42.3
8 B	323	1.39	61058	17.2763	2.7	0.70464	3.9	0.088331	2.8	0.712	-0.67	545.7	14.6	541.6	16.4	524.3	60.1
12 B	867	0.77	488045	17.0969	2.3	0.72821	3.2	0.090338	2.2	0.68	-0.34	557.5	11.6	555.5	13.6	547.2	50.9
10 B	564	1.69	1286746	17.1749	3	0.71118	5.1	0.088627	4.1	0.808	-0.3	547.4	21.7	545.4	21.6	537.2	66.1
10 A	407	1.82	589720	17.0432	1.9	0.74566	3.4	0.092211	2.9	0.83	-0.41	568.6	15.6	565.7	15	554	42
12 A	897	0.73	239259	16.9237	1.9	0.72135	2.9	0.08858	2.3	0.774	0.67	547.1	11.9	551.5	12.5	569.4	40.4
12 D	719	0.86	74762	17.1815	2.2	0.71074	3.2	0.088607	2.3	0.736	-0.35	547.3	12.3	545.2	13.5	536.4	47.3
8 C	310	1.38	22655	17.1563	2.6	0.72132	4	0.089793	3.1	0.765	-0.44	554.3	16.4	551.4	17.1	539.6	56.8
13 B	295	1.25	174812	17.1984	2.5	0.75066	4	0.093676	3.1	0.782	-1.3	577.2	17.2	568.6	17.3	534.2	54.1
13 A	472	1.1	61888	17.1584	2.4	0.73705	3.9	0.091764	3.1	0.79	-0.77	566	16.5	560.7	16.6	539.4	51.8
14 D	413	1.57	145640	17.3853	2.3	0.7176	3.2	0.090523	2.1	0.679	-1.6	558.6	11.5	549.2	13.4	510.5	51
14 B*	521	1.85	37874	16.2042	3.6	0.8031	4.6	0.094426	2.9	0.617	2.8	581.7	15.9	598.6	21	663.2	78.2
14 A	440	1.72	18688	17.482	2.5	0.71095	3.9	0.090183	3	0.766	-1.7	556.6	16.1	545.3	16.6	498.3	55.7
15 A	400	2.3	419812	16.955	2.7	0.70894	4.4	0.087217	3.5	0.788	0.78	539.1	18	544.1	18.6	565.4	59.4
6 D	480	0.91	53177	17.2256	1.8	0.71525	3.4	0.089397	2.9	0.842	-0.6	552	15.1	547.9	14.4	530.8	40
551063																	
1 R	113	1.63	18347	17.3389	3.3	0.7062	5	0.088847	3.7	0.747	-1	548.7	19.6	542.5	21	516.4	73
3 R	69	1.88	19452	17.384	4.1	0.72339	6.3	0.091247	4.8	0.764	-1.6	562.9	25.9	552.7	26.8	510.7	89.2
3 C	149	1.17	6828	17.5172	6.3	0.68754	7.9	0.087389	4.7	0.598	-1.6	540.1	24.6	531.3	32.8	493.9	139.9
6 R	87	1.31	14491	17.187	3.3	0.71991	5.5	0.089779	4.4	0.797	-0.54	554.2	23.2	550.6	23.3	535.7	72.6
8*	114	2.15	24997	17.3059	4	0.73888	4.9	0.092782	2.7	0.565	-1.8	572	15	561.8	21	520.6	87.9
11 C	134	1.38	9723	17.4824	4.2	0.69571	5.4	0.088251	3.4	0.632	-1.5	545.2	17.8	536.2	22.5	498.3	92.1
13	244	1.23	24668	17.1319	2.7	0.72322	3.7	0.089902	2.5	0.671	-0.4	555	13.1	552.6	15.6	542.7	59.5
14	158	1.29	24304	17.4692	3.3	0.70396	4.6	0.089231	3.2	0.704	-1.6	551	17.1	541.2	19.3	500	71.9
16 R	266	1.2	13381	17.1193	3.1	0.71175	5.4	0.088412	4.3	0.812	-0.047	546.1	22.7	545.8	22.6	544.3	68.2
17	136	1.71	14890	16.9466	3.8	0.73443	5.3	0.090309	3.7	0.697	0.29	557.4	19.6	559.2	22.6	566.5	82.2
18 C	95	2.03	7970	17.4356	5.4	0.72274	6.7	0.091436	3.9	0.584	-2	564	21.1	552.3	28.5	504.2	119.4
18 R	228	2.35	22653	17.1076	3.1	0.71115	4.6	0.088276	3.3	0.73	0.017	545.3	17.5	545.4	19.3	545.8	68.5
20 C	121	1.7	6793	17.0323	5.5	0.72722	6.5	0.089874	3.5	0.533	0.025	554.8	18.4	554.9	27.7	555.5	119.6
20 R	81	2.07	39407	16.4472	4.7	0.74348	6.4	0.088727	4.3	0.669	2.8	548	22.5	564.4	27.7	631.2	102.3
23	101	1.3	52768	17.0676	4.2	0.70869	5.9	0.087765	4.1	0.703	0.28	542.3	21.4	544	24.7	550.9	91
24	205	1.59	28539	17.1556	2.5	0.71011	3.8	0.088395	2.9	0.754	-0.2	546	15	544.8	16	539.7	

Table 1. LA-ICP-MS U-Pb zircon data for the Southern Pines 100 K (crystallization ages)

Composition			Isotopic Ratios									Ages (Ma)					
ID	U (ppm)	U/Th	²⁰⁶ Pb/ ²⁰⁴ Pb	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (%)	²⁰⁷ Pb/ ²³⁵ U	Error (%)	²⁰⁶ Pb/ ²³⁸ U	Error (%)	error correlation	concordia distance (d _c) ^a	²⁰⁶ Pb/ ²³⁸ U	Error (Ma)	²⁰⁷ Pb/ ²³⁵ U	Error (Ma)	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (Ma)
25	58	1.64	20477	16.7764	6.2	0.70336	8	0.085619	5	0.631	2	529.6	25.7	540.8	33.6	588.4	134.7
26	149	2.37	20567	17.5752	3.7	0.65252	5.2	0.083213	3.7	0.714	-0.91	515.3	18.5	510	21	486.6	81.2
27	152	3.14	37090	17.4951	3.5	0.69796	5.3	0.088601	4	0.754	-1.5	547.3	20.9	537.6	22.1	496.7	76.5
28*	485	1.37	1127079	17.0919	2.2	0.74746	3.7	0.092698	3	0.802	-0.68	571.5	16.1	566.7	16	547.8	48
29	220	1.48	28149	17.2024	3.4	0.71201	4.8	0.088872	3.4	0.707	-0.48	548.9	17.8	545.9	20.2	533.8	74.1
7 C	123	1.48	40803	17.1169	3.8	0.72433	5.4	0.089961	3.8	0.702	-0.34	555.3	20.1	553.2	22.9	544.6	83.6
31	76	1.71	56752	17.3321	5	0.70401	7.1	0.088537	5	0.714	-0.94	546.9	26.5	541.2	29.7	517.2	108.7
32	272	1.35	77384	17.2151	2.7	0.70479	3.8	0.088036	2.7	0.709	-0.38	543.9	14.1	541.6	15.9	532.1	58.6
33	316	1.63	66010	17.107	3	0.69032	3.8	0.085687	2.4	0.635	0.53	530	12.4	533	15.9	545.9	64.7
12 C	188	1.16	162653	17.1769	3.9	0.70189	6	0.087479	4.6	0.756	-0.11	540.6	23.6	539.9	25.2	537	86.4
35	214	1.51	106587	16.952	3.5	0.69381	5.7	0.085341	4.5	0.789	1.1	527.9	22.7	535.1	23.7	565.8	76.1
37	319	1.18	18115	17.3556	4	0.6893	4.8	0.086804	2.7	0.558	-0.76	536.6	13.9	532.4	20	514.3	88
38	204	1.67	58994	17.007	4.2	0.69894	6.4	0.086251	4.8	0.748	0.79	533.3	24.4	538.2	26.6	558.7	92.1
39	113	2.36	104950	17.3438	4.1	0.72549	5.7	0.0913	4	0.705	-1.5	563.2	21.7	553.9	24.4	515.8	89
40	89	1.91	7067	17.801	6.6	0.64462	7.7	0.083261	3.9	0.515	-2	515.6	19.5	505.2	30.5	458.3	145.6
41	254	1.78	49251	17.2972	3.7	0.68025	5	0.085377	3.4	0.668	-0.21	528.1	17	526.9	20.7	521.7	82.1
42	195	1.63	26180	17.5525	2.9	0.69041	4.2	0.087931	3	0.716	-1.7	543.3	15.7	533	17.5	489.5	65
43*	573	13.25	66476	16.8498	2.3	0.76324	5.2	0.093314	4.6	0.895	0.11	575.1	25.4	575.9	22.6	578.9	50
44 C	532	1.19	70375	17.2618	2.3	0.72022	4.4	0.090208	3.7	0.846	-0.87	556.8	19.8	550.8	18.7	526.2	51.4
44 R	441	1.28	31693	17.3674	2.9	0.72104	4.4	0.090863	3.3	0.751	-1.4	560.6	17.6	551.3	18.6	512.8	63.2

Notes:

^aConcordia distance (d_c) was calculated using IsoplotR (Vermeesch, 2018), and we discarded analyses outside of the range -2 < d_c < 4.6 (Vermeesch, 2021).

*Analyses with an asterisk were considered outliers according to Chauvenet’s Criterion in IsoplotR (Vermeesch, 2018). They were not included in the weighted mean.

Table 2. LA-ICP-MS U-Pb zircon data for the Southern Pines 100 K (detrital ages)

Composition			Isotopic Ratios								Ages (Ma)						
ID	U (ppm)	Th/U	²⁰⁶ Pb/ ²⁰⁴ Pb	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (%)	²⁰⁷ Pb/ ²³⁵ U	Error (%)	²⁰⁶ Pb/ ²³⁸ U	Error (%)	error correlation	concordia distance (d _c) ^a	²⁰⁶ Pb/ ²³⁸ U	Error (Ma)	²⁰⁷ Pb/ ²³⁵ U	Error (Ma)	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (Ma)
Southern Pines 100k																	
Sample 551983																	
1	170	0.43	586561	10.3823	1.8	3.51978	3.9	0.265157	3.5	0.892	1.3	1516.2	46.7	1531.7	30.7	1553.1	32.9
2	40	0.32	18242	12.7737	5.1	2.15128	7.7	0.199393	5.8	0.751	-0.5	1172.1	62.2	1165.5	53.6	1153.2	101.3
3	79	0.50	26612	11.6707	3.3	2.60115	5.4	0.22027	4.3	0.795	1.2	1283.3	49.7	1301	39.4	1330.2	63.1
4	376	0.96	166729	17.1558	3	0.73002	6	0.090874	5.2	0.871	-0.6	560.7	28.1	556.6	25.8	539.7	64.7
5	343	0.31	238147	12.3697	2.1	2.3429	3.8	0.210284	3.2	0.834	-0.38	1230.3	35.8	1225.4	27.2	1216.7	41.6
6	120	0.57	199579	17.1206	5	0.69874	6	0.086802	3.4	0.564	0.26	536.6	17.4	538	25.1	544.2	108.3
11	176	0.93	34688	16.8935	3.6	0.70179	5	0.086025	3.5	0.697	1.3	532	17.7	539.9	20.8	573.3	77.7
12	267	0.20	111479	12.502	1.8	2.32514	3.8	0.210922	3.3	0.879	-1.3	1233.7	37.2	1220	26.7	1195.8	35.3
13	92	0.88	28406	16.7306	5.1	0.73435	7.3	0.089147	5.2	0.716	1.4	550.5	27.4	559.1	31.2	594.3	109.8
16	144	0.52	157949	10.796	2.3	3.10217	5.4	0.243009	4.9	0.903	2.8	1402.3	61.6	1433.2	41.5	1479.4	44
17	147	1.16	303295	16.7932	3.3	0.70824	6.6	0.0863	5.7	0.862	1.5	533.6	28.9	543.7	27.6	586.2	72.2
18	165	0.79	258172	8.8401	3.6	5.31268	5.7	0.340772	4.4	0.772	-1	1890.4	72.1	1870.9	48.7	1849.3	65.5
19	3033	0.60	103264	16.5531	2.7	0.73822	3.5	0.088666	2.1	0.61	2.4	547.6	11.1	561.4	14.9	617.4	59.3
21	242	0.91	123296	16.5853	4	0.71202	5.7	0.085686	4	0.71	2.8	530	20.4	545.9	23.9	613.2	86.1
22	186	0.67	26190	16.8651	3.8	0.71306	5.9	0.087258	4.5	0.764	1.2	539.3	23.2	546.6	24.8	576.9	82.3
23	123	0.65	31504	17.1908	3.4	0.69846	6.2	0.087123	5.2	0.836	-0.092	538.5	26.8	537.9	25.9	535.2	74.5
24	133	1.03	60842	16.9939	4.6	0.70928	9.5	0.087459	8.4	0.878	0.57	540.5	43.4	544.3	40.1	560.4	99.3
26	64	0.71	10737	17.2783	5.9	0.6604	13.1	0.082795	11.7	0.892	0.34	512.8	57.7	514.9	53	524.1	129.8
27	623	0.33	153575	16.3774	1.8	0.88189	3.3	0.104797	2.7	0.828	-0.061	642.5	16.6	642	15.6	640.4	39.5
28	249	0.53	276346	16.3614	3.1	0.83508	4.3	0.099139	3	0.7	1.1	609.4	17.5	616.4	19.8	642.5	65.8
29	169	0.90	33322	17.0955	4.7	0.69816	6.1	0.086603	3.8	0.631	0.41	535.4	19.7	537.7	25.4	547.4	103.2
30	24	0.45	28970	16.0426	9.1	0.88815	13.4	0.103384	9.9	0.738	1.5	634.2	59.9	645.4	64.1	684.7	193.3
31	111	0.58	72173	16.59	5.5	0.70712	6.6	0.08512	3.5	0.535	3.1	526.6	17.8	543	27.6	612.5	119.7
32	75	0.55	37952	9.2202	2.4	4.55806	6.9	0.304939	6.4	0.937	2.4	1715.8	97	1741.6	57.2	1772.8	43.7
33	376	0.95	699888	9.9182	1.4	4.0389	3.6	0.290662	3.3	0.921	-0.25	1644.8	47.5	1642.1	28.9	1638.5	25.7
34	48	0.84	10489	17.1024	8.3	0.69959	10.2	0.086816	6	0.587	0.33	536.7	30.9	538.5	42.7	546.5	180.8
35	191	0.88	70832	16.4251	3.3	0.86837	5.3	0.103492	4.1	0.786	-0.02	634.8	25	634.7	24.9	634.1	70.1
39	120	1.22	93573	16.3817	5.2	0.69249	7.4	0.082313	5.3	0.719	4.4	509.9	26.1	534.3	30.8	639.8	110.8
40	835	0.22	171288	15.1125	2.2	1.12861	3.8	0.123758	3	0.806	1.7	752.2	21.5	767.1	20.2	810.8	46.5
41	110	0.47	101257	16.757	4.2	0.74458	6.9	0.090532	5.5	0.795	0.96	558.7	29.3	565.1	29.8	590.9	90.6
43	561	0.19	197967	12.287	1.5	2.3833	3.2	0.21248	2.8	0.886	-0.42	1242	31.7	1237.6	22.6	1229.9	28.8
44	131	0.40	27587	16.0283	5.1	0.87985	7	0.102327	4.8	0.686	1.9	628	28.6	640.9	33.1	686.5	107.9
45	44	0.29	73444	9.4276	3.5	4.41426	7	0.301963	6.1	0.869	1	1701	91	1715	58	1732.1	63.7
46	242	0.51	65783	16.6949	2.8	0.78832	5.2	0.095495	4.4	0.846	0.31	588	24.6	590.2	23.1	598.9	59.6
47	94	0.71	142985	10.2596	2	3.68763	4.9	0.274519	4.5	0.914	0.45	1563.7	62.6	1568.7	39.4	1575.4	37.6
48	154	0.93	778376	17.0351	3.8	0.69899	6.6	0.0864	5.4	0.818	0.61	534.2	27.5	538.2	27.4	555.1	82.2
50	122	0.48	23903	16.0153	3.8	0.87278	6.3	0.101422	5	0.791	2	622.7	29.6	637.1	29.8	688.3	82
52	134	0.56	25734	16.8051	5.1	0.7452	6.7	0.090868	4.4	0.655	0.8	560.7	23.6	565.4	29.1	584.7	110.2
53	114	0.68	50355	16.1176	6	0.93167	7.6	0.108958	4.6	0.61	0.27	666.7	29.3	668.5	37.2	674.6	128.6
54	334	0.68	402385	16.6923	2.2	0.72606	3.4	0.087939	2.6	0.762	1.7	543.3	13.4	554.2	14.4	599.3	47.2
55	93	0.79	27722	17.1349	3.8	0.70611	7	0.087791	5.9	0.839	-0.003	542.5	30.7	542.4	29.5	542.3	83.7
56	185	0.88	196802	16.2387	4	0.83213	5.1	0.098048	3.3	0.637	1.9	603	18.9	614.8	23.7	658.6	85.1
57	551	0.64	550231	16.3775	1.9	0.84306	3.8	0.100185	3.3	0.869	0.71	615.5	19.5	620.8	17.8	640.4	40.8
58	135	0.95	73918	15.748	5	0.95186	8.6	0.108766	6.9	0.81	1.7	665.6	43.9	679.1	42.5	724	106.7
60	57	0.87	23872	17.0475	6.2	0.72359	8.8	0.089505	6.2	0.707	0.022	552.6	33	552.8	37.6	553.5	136.1
61	67	0.74	19786	16.722	6	0.70647	7.6	0.085719	4.7	0.622	2.2	530.2	24.1	542.7	32	595.4	129.2
62	263	0.38	253625	8.6012	1.8	5.30361	3.8	0.330998	3.3	0.875	1.8	1843.2	53.1	1869.4	32.4	1898.7	33
64																	

Table 2. LA-ICP-MS U-Pb zircon data for the Southern Pines 100 K (detrital ages)

Composition			Isotopic Ratios								Ages (Ma)						
ID	U (ppm)	Th/U	²⁰⁶ Pb/ ²⁰⁴ Pb	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (%)	²⁰⁷ Pb/ ²³⁵ U	Error (%)	²⁰⁶ Pb/ ²³⁸ U	Error (%)	error correlation	concordia distance (d _c) ^a	²⁰⁶ Pb/ ²³⁸ U	Error (Ma)	²⁰⁷ Pb/ ²³⁵ U	Error (Ma)	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (Ma)
136	92	0.23	605367	11.6132	3.5	2.64527	5.6	0.222903	4.4	0.781	1.1	1297.2	51.4	1313.3	41.3	1339.7	67.7
137	41	0.74	236148	16.4229	5.5	0.83148	8.8	0.099082	6.8	0.782	0.76	609	39.8	614.4	40.4	634.4	117.4
138	176	1.01	154125	16.0101	3.3	0.8918	5.3	0.103599	4.2	0.791	1.6	635.5	25.4	647.3	25.4	688.9	69.3
139	139	0.43	18331	16.4116	4.9	0.7991	6.1	0.095158	3.6	0.591	1.7	586	20.1	596.3	27.4	635.9	105.6
140	173	1.19	72013	17.2744	4.5	0.69431	6	0.087027	4	0.665	-0.44	537.9	20.7	535.4	25.1	524.6	98.9
142	283	0.32	471811	15.9912	2.5	0.89992	3.6	0.104419	2.6	0.719	1.6	640.3	15.7	651.7	17.2	691.5	53
143	190	0.28	38774	16.482	3.5	0.79071	4.6	0.094563	2.9	0.643	1.5	582.5	16.4	591.6	20.5	626.6	75.5
144	200	0.56	490498	6.2053	2	10.51143	4.7	0.47328	4.3	0.904	-0.96	2497.9	88.4	2480.9	43.8	2467	34
145	434	0.16	403891	12.0045	3.2	2.4273	3.9	0.211427	2.1	0.547	1.3	1236.4	23.8	1250.7	27.8	1275.4	63.1
146	269	0.83	29956	17.3132	3.4	0.68951	4.9	0.086619	3.6	0.734	-0.49	535.5	18.6	532.5	20.5	519.6	73.7
147	247	1.02	106864	16.6066	3.1	0.80109	5.2	0.096529	4.2	0.809	0.47	594	24.1	597.4	23.7	610.4	66.7
148	333	0.49	1877577	16.3046	2.9	0.86773	4.1	0.102657	2.9	0.702	0.63	630	17.4	634.3	19.5	650	63.1
149	77	1.37	244136	16.9628	5.5	0.72149	8.3	0.088802	6.2	0.744	0.5	548.4	32.5	551.5	35.3	564.3	120.8
150	85	0.62	17548	16.565	6.2	0.72761	7.4	0.087455	4.1	0.549	2.7	540.5	21.2	555.2	31.8	615.8	134.3
151	48	0.64	132928	16.6906	6	0.71057	8.7	0.086054	6.3	0.721	2.2	532.2	32	545.1	36.6	599.5	130.4
152	140	0.85	46850	16.7072	4.6	0.72452	8.1	0.087831	6.7	0.822	1.6	542.7	34.8	553.3	34.7	597.3	100.5
153	124	0.54	246819	15.6725	3.9	1.02199	5.4	0.11622	3.6	0.677	0.83	708.8	24.3	714.9	27.5	734.3	83.4
154	202	0.23	92368	13.701	2.4	1.69984	3.6	0.168988	2.6	0.741	0.17	1006.5	24.7	1008.5	22.8	1012.7	48.5
156	175	0.69	76933	16.9446	3.8	0.736	5.7	0.090491	4.3	0.751	0.25	558.4	23.1	560.1	24.8	566.7	82.6
157	130	0.98	33645	16.7419	5	0.72157	6.3	0.087655	3.9	0.613	1.7	541.7	20.2	551.6	27	592.9	108.7
158	109	0.51	110178	9.8829	2.3	3.80772	4.9	0.273052	4.4	0.89	3.1	1556.3	60.8	1594.4	39.7	1645.1	41.8
160	368	0.12	638714	11.6564	2.1	2.64657	4	0.223843	3.3	0.842	0.86	1302.1	39.2	1313.7	29.1	1332.6	41.2
161	63	0.73	38103	16.7188	5.5	0.72037	8.7	0.087389	6.8	0.778	1.7	540.1	35	550.9	36.9	595.8	118.3
162	125	1.19	15230	15.9602	3.8	0.8589	5.5	0.099466	4	0.726	2.7	611.3	23.4	629.5	25.9	695.6	81
163	117	0.67	83832	16.812	5.3	0.72112	6.3	0.087967	3.4	0.541	1.4	543.5	17.9	551.3	27	583.8	115.8
164	64	0.81	71854	16.6341	4.2	0.7224	6	0.08719	4.3	0.718	2.2	538.9	22.3	552.1	25.6	606.8	90.6
165	75	0.78	116417	16.5888	4.6	0.81303	7.7	0.097862	6.2	0.803	0.32	601.9	35.7	604.2	35.2	612.7	99.4
166	116	0.62	117464	16.7777	3.9	0.73277	6	0.089206	4.6	0.76	1.1	550.8	24.3	558.2	25.9	588.2	85.1
167	1112	0.96	227360	16.3961	1.7	0.82578	2.5	0.098242	1.8	0.71	1.1	604.1	10.1	611.3	11.3	637.9	37.4
168	209	0.81	51240	16.5155	2.9	0.83846	4.4	0.100478	3.3	0.757	0.15	617.2	19.5	618.3	20.3	622.3	61.7
169	138	0.76	48912	17.0042	4.5	0.70496	5.3	0.086979	2.8	0.534	0.75	537.6	14.6	541.7	22.2	559.1	97.5
170	199	0.88	72471	17.1336	3.9	0.7135	5.4	0.088703	3.7	0.697	-0.17	547.9	19.7	546.8	22.7	542.5	84.2
171	147	0.72	55638	16.892	3.8	0.6969	5.5	0.085418	4	0.722	1.4	528.4	20.1	536.9	22.9	573.5	82.7
172	197	0.77	89507	16.4426	3.9	0.75817	5.1	0.090454	3.4	0.661	2.5	558.2	18.2	573	22.5	631.8	83
174	86	0.63	10955	16.7838	5.9	0.71144	7.6	0.086641	4.8	0.629	1.8	535.6	24.6	545.6	32.1	587.4	128.3
175	211	0.72	67644	16.0127	4	0.83681	5.1	0.097227	3.2	0.627	3.1	598.1	18.4	617.4	23.8	688.6	85.4
176	154	0.96	28446	17.1183	4.3	0.71521	6	0.088836	4.1	0.692	-0.14	548.6	21.7	547.8	25.2	544.5	93.9
178	461	0.30	170418	10.4922	1.5	3.49703	3.9	0.266231	3.6	0.924	0.47	1521.6	49.1	1526.5	31	1533.3	28.3
180	147	0.35	59900	17.0067	4.3	0.75548	6.4	0.093226	4.7	0.735	-0.49	574.6	25.8	571.4	27.9	558.8	94.3
182	74	0.92	61646	16.6297	6.4	0.72847	8	0.0879	4.8	0.598	2.2	543.1	25.1	555.7	34.4	607.4	139.4
183	227	0.68	99220	17.0792	3	0.70337	4.7	0.087166	3.7	0.78	0.32	538.8	19.1	540.8	19.9	549.5	64.9
184	126	0.70	23617	17.0188	3.5	0.77811	5.9	0.096087	4.8	0.813	-0.98	591.4	27.3	584.4	26.4	557.2	75.5
185	75	0.72	69945	15.8187	5.3	0.896	7.4	0.102842	5.3	0.709	2.7	631	31.7	649.6	35.7	714.6	111.6
186	369	0.72	258134	16.7353	1.9	0.79307	3.4	0.096302	2.9	0.832	0.03	592.7	16.1	592.9	15.4	593.7	41.2
187	113	0.73	35106	16.8082	4	0.69686	5.9	0.084989	4.4	0.739	1.9	525.8	22.2	536.9	24.8	584.3	87
188	203	1.02	88697	16.6054	3.2	0.75139	6.6	0.090534	5.7	0.875	1.5	558.7	30.7	569	28.6	610.5	68.7
189	192	0.48	229304	9.7877	2	4.11503	3.8	0.292246	3.2	0.843	0.3	1652.7	46.2	1657.3	30.7	1663	37.4
190	57	0.81	15634	16.7929	5.3	0.73736	7.5	0.089846	5.3	0.708	1	554.6	28.2	560.9	32.2	586.2	114.7
191	424	0.45	166174	16.4645	2	0.86812	2.9	0.103711	2	0.707	-0.23	636.1	12.3	634.5	13.6	628.9	43.8
194	95	0.56	12888	17.2012	4.8	0.71313	7.4	0.089007	5.6	0.765	-0.49	549.7	29.7	546.6	31.2	533.9	104.1
195	66	0.58	166758	9.3055	2.6	4.6023	6.1	0.310748	5.5	0.9	0.41	1744.4	83.5	1749.7	50.6	1756	48.3
197	171	0.60	210790	17.036	3.7	0.73722	4.6	0.09113	2.7	0.593	-0.25	562.2	14.7	560.8	19.9	555	81.2
198	213	0.88	199749	16.5447	3.4	0.86089	4.9	0.103348	3.5	0.721	-0.48	634	21.4	630.6	23.1	618.5	73.5
199	427	0.88	1743713	16.9361	2.8	0.72671	4.2	0.089303	3.1	0.741	0.52	551.4	16.4	554.6	17.9	567.8	61.4
200	403	0.31	3938376	13.7254	1.8	1.69812	3.9	0.169117	3.4	0.887	0.059	1007.2	31.9	1007.8	24.6	1009.1	36
202	272	1.12	52630	16.6025	2.7	0.82753	4.2	0.099691	3.2	0.771	-0.05	612.6	18.8	612.2	19.2	610.9	57.5
203	34	0.60	16049	17.4132	8.2	0.70021	10.7	0.08847	6.9	0.646	-1.3	546.5	36.2	538.9	44.8	507	179.9
205	80	1.12	10725	4.6249	5.4	16.92998	8.8	0.568134	7	0.793	1.2	2900.2	163.4	2930.9	84.6	2952	86.6
207	123	0.89	602942	16.6627	4.3	0.71304	6.8	0.08621	5.2	0.772	2.2	533.1	26.7	546.6	28.6	603.1	93
208	94	0.97	22978	16.8624	5.6	0.74264	7.3	0.090864	4.6	0.637	0.57	560.6	24.9	563.9	31.5	577.3	122
209	167	0.66	27860	17.061	4.8	0.71032	6	0.087933	3.6	0.601	0.29	543.3	18.7	544.9	25.2	551.8	104.1
211	88	0.40	205386	16.3392	5.6	0.80933	8.2	0.095951	5.9	0.724	1.8	590.6	33.4	602.1	37.1	645.4	121.1
212	117	0.75	20342	17.2022	4.1	0.71035	6.3	0.088664	4.9	0.765	-0.41	547.6	25.5	545	26.7	533.8	89.4
213	102	0.41	30158	16.7134	4.4	0.74561	6.5	0.090421	4.8	0.739	1.2	558	25.6	565.7	28.1	596.5	94.6
214	196	0.83	26871	16.0375	2.7	0.94853	4.1	0.110377	3.1	0.758	0.31	674.9	19.9	677.3	20.3	685.3	57.1
215	491	0.61	259933	16.522	1.9	0.83554	3.8	0.100167	3.3	0.873	0.17	615.4	19.4	616.7	17.5	621.4	40
216	254	0.76	73892	16.9861	3.4	0.75784	4.5	0.093403	2.9	0.651	-0.47	575.6	16.2	572.8	19.8	561.4	74.7
218	138	0.25	150584	12.4119	2.2	2.2714	5.2	0.204563	4.7	0.903	0.38	1199.8	50.9	1203.4	36.3	1210	43.6

Table 2. LA-ICP-MS U-Pb zircon data for the Southern Pines 100 K (detrital ages)

Composition			Isotopic Ratios									Ages (Ma)					
ID	U (ppm)	Th/U	²⁰⁶ Pb/ ²⁰⁴ Pb	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (%)	²⁰⁷ Pb/ ²³⁵ U	Error (%)	²⁰⁶ Pb/ ²³⁸ U	Error (%)	error correlation	concordia distance (d _c) ^a	²⁰⁶ Pb/ ²³⁸ U	Error (Ma)	²⁰⁷ Pb/ ²³⁵ U	Error (Ma)	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (Ma)
263	200	0.44	27639	18.152	4.5	0.50828	5.8	0.066945	3.7	0.634	-0.1	417.7	14.8	417.3	19.8	414.9	100
265	111	0.61	26487	16.3468	5	0.728	6.7	0.086349	4.5	0.668	3.8	533.9	23.1	555.4	28.8	644.4	107.8
266	217	0.39	123515	16.2743	3.1	0.85818	4.2	0.101339	2.8	0.665	1	622.2	16.5	629.1	19.6	653.9	67.1
267	405	0.47	6115573	10.5285	1.6	3.43769	3.7	0.262619	3.3	0.898	0.86	1503.2	44.8	1513.1	29.3	1526.8	30.9
269	74	0.81	212213	16.6354	5	0.7255	7	0.087572	4.9	0.695	2.2	541.2	25.2	553.9	29.8	606.6	108.6
270	171	0.71	153350	16.5508	3.7	0.85889	5.8	0.103145	4.4	0.758	-0.45	632.8	26.3	629.5	27	617.7	80.9
271	393	0.33	386389	10.3222	1.9	3.72835	3.5	0.279243	3	0.844	-0.69	1587.5	41.8	1577.5	28.2	1564	35.3
273	130	0.26	83114	11.194	3	3.05944	5.7	0.248496	4.8	0.846	-0.6	1430.7	61.5	1422.6	43.4	1410.4	57.9
274	182	0.31	52540	16.1589	3.1	0.82617	5.4	0.096867	4.4	0.813	2.1	596	24.9	611.5	24.7	669.2	67
277	66	0.49	35798	17.395	7.2	0.55271	8.1	0.069761	3.8	0.468	2.7	434.7	16	446.8	29.4	509.3	158.1
278	118	0.59	22879	17.108	5.2	0.6953	6.7	0.086311	4.3	0.64	0.41	533.7	22.1	536	28.1	545.7	113.2
280	157	0.67	77433	16.9114	4.1	0.7485	5.6	0.091847	3.7	0.671	0.15	566.5	20.3	567.4	24.3	571	90
286	272	0.87	164524	16.234	3.9	0.86165	4.5	0.101496	2.2	0.502	1.3	623.2	13.3	631	20.9	659.3	82.7
287	177	0.61	30772	16.6367	4.2	0.73566	5.8	0.088806	4	0.684	1.9	548.5	20.9	559.9	25	606.5	91.6
288	138	0.48	186616	7.3651	1.7	7.56382	3.6	0.404219	3.2	0.878	-0.48	2188.4	59.4	2180.6	32.7	2173.2	30.4
289	37	0.89	54353	7.3404	2.7	7.27643	6	0.387556	5.4	0.892	2.2	2111.5	96.6	2145.9	53.7	2179	47.3
290	131	1.27	680416	16.4815	4.3	0.76264	6.3	0.091203	4.7	0.737	2	562.7	25.1	575.5	27.7	626.7	91.9
291	159	0.18	299555	16.1719	2.8	0.84328	4.3	0.098953	3.2	0.75	1.9	608.3	18.6	621	19.8	667.5	60.4
292	184	0.09	719248	5.489	1.9	12.50029	3.9	0.497857	3.4	0.877	2	2604.6	72.7	2642.7	36.4	2672.1	30.8
293	150	0.75	107459	17.0318	3	0.70296	5.4	0.086872	4.5	0.826	0.54	537	23	540.6	22.7	555.5	66.5
294	357	0.53	309847	17.0362	2.7	0.70513	3.2	0.087163	1.6	0.515	0.58	538.7	8.4	541.8	13.3	555	59.3
295	127	0.99	61427	16.1754	3	0.84901	4.7	0.099647	3.5	0.758	1.7	612.3	20.7	624.1	21.8	667	65.3
296	93	0.45	1217845	10.7406	2.6	3.34285	4.7	0.260519	3.9	0.825	-0.094	1492.5	51.5	1491.1	36.6	1489.2	50.1
297	69	0.70	20642	16.8479	6.1	0.73574	8.6	0.089942	6.1	0.706	0.77	555.2	32.4	559.9	37.1	579.1	132.8
298	362	0.87	2093740	16.3422	2.5	0.85225	4.1	0.101058	3.3	0.791	0.71	620.6	19.3	625.9	19.2	645	54.1
300	399	0.72	151091	16.9741	2.7	0.70769	3.9	0.087162	2.9	0.734	0.76	538.7	14.9	543.4	16.5	562.9	57.9
301	203	0.30	184832	10.5553	2.2	3.42304	3.9	0.262166	3.2	0.824	0.58	1500.9	43	1509.7	30.6	1522	41.6
302	271	0.30	556351	10.4223	2	3.4929	4.1	0.264147	3.6	0.872	1.1	1511	48	1525.6	32.2	1545.9	37.5
303	122	0.99	28654	16.951	4.8	0.74544	6.3	0.091685	4.1	0.648	0.0096	565.5	22.1	565.6	27.4	565.9	104.8
304	268	0.67	44133	16.7314	3.2	0.73733	4.7	0.089514	3.4	0.731	1.3	552.7	18.1	560.8	20.1	594.2	68.9
305	84	0.66	38614	16.8001	6.3	0.68982	8.1	0.084089	5.1	0.627	2.2	520.5	25.4	532.7	33.5	585.3	136.7
307	94	0.25	55341	16.4575	4.4	0.75604	6.4	0.090282	4.6	0.722	2.4	557.2	24.7	571.7	28	629.9	95.6
308	117	1.41	17826	16.9328	5	0.75884	6.7	0.093233	4.4	0.664	-0.21	574.6	24.4	573.3	29.3	568.2	108.9
309	163	0.85	575098	16.3037	4.8	0.72653	6.3	0.085947	4.1	0.647	4.1	531.5	20.9	554.5	27	650.1	103.6
310	128	0.94	187807	17.1279	3.9	0.70489	5.4	0.087603	3.8	0.697	0.063	541.3	19.7	541.7	22.9	543.2	85.4
313	43	1.00	16698	16.584	6	0.71227	8.6	0.085709	6.1	0.716	2.8	530.1	31.2	546.1	36.2	613.3	129.1
315	194	0.29	57478	11.752	2.3	2.5674	4.2	0.218927	3.5	0.833	1.1	1276.2	40.6	1291.4	30.7	1316.7	45.1
<i>Sample 551641</i>																	
2	109	0.38	99976	17.3845	5.2	0.70866	8.6	0.08939	6.9	0.798	-1.2	551.9	36.5	543.9	36.4	510.7	114.4
3	268	0.10	68307	15.903	3.4	0.95136	4.8	0.109779	3.3	0.693	1	671.5	21.1	678.8	23.6	703.3	73.1
5	855	0.04	56410	18.4744	2.6	0.44608	4.2	0.059797	3.3	0.781	0.03	374.4	11.9	374.5	13.1	375.4	58.8
6	304	0.41	116210	13.3644	2.3	1.83717	4.7	0.178152	4.1	0.871	0.19	1056.9	39.9	1058.8	30.9	1062.9	46.5
9	518	0.01	74023	17.8681	5.7	0.48222	8.1	0.062521	5.7	0.709	2	390.9	21.8	399.6	26.7	450	126.8
11	360	0.56	447419	17.0763	2.7	0.73279	3.9	0.090796	2.8	0.72	-0.33	560.2	15.1	558.2	16.8	549.8	59.4
13	279	0.60	91634	16.3064	3	0.82427	5.8	0.097527	5	0.858	1.4	599.9	28.6	610.4	26.7	649.7	64.2
14	817	0.04	53525	18.2707	2.4	0.47309	3.6	0.062718	2.7	0.746	0.26	392.1	10.1	393.3	11.6	400.3	53
18	240	0.26	195909	10.5439	2	3.41542	4.5	0.2613	4	0.9	1	1496.5	53.9	1507.9	35.2	1524.1	36.8
19	211	0.48															

Table 2. LA-ICP-MS U-Pb zircon data for the Southern Pines 100 K (detrital ages)

Composition			Isotopic Ratios								Ages (Ma)						
ID	U (ppm)	Th/U	²⁰⁶ Pb/ ²⁰⁴ Pb	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (%)	²⁰⁷ Pb/ ²³⁵ U	Error (%)	²⁰⁶ Pb/ ²³⁸ U	Error (%)	error correlation	concordia distance (d _c) ^a	²⁰⁶ Pb/ ²³⁸ U	Error (Ma)	²⁰⁷ Pb/ ²³⁵ U	Error (Ma)	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (Ma)
22	793	0.79	67404	17.3941	2.9	0.69424	3.7	0.087621	2.4	0.639	-1.1	541.5	12.4	535.3	15.6	509.4	63.3
23	278	0.44	108603	10.6132	1.9	3.35177	4	0.258117	3.5	0.875	1.1	1480.2	46.4	1493.2	31.4	1511.7	36.6
24	106	0.48	21181	16.8181	5.8	0.81357	7.4	0.099281	4.7	0.629	-0.89	610.2	27.1	604.5	33.8	583	125.1
26	123	0.68	16468	17.4181	4	0.69614	6.7	0.087981	5.4	0.802	-1.1	543.6	28.1	536.5	28	506.4	88.3
28	99	0.79	19640	17.2105	4.6	0.73806	7.6	0.092168	6.1	0.794	-1	568.3	33	561.3	33	532.7	101.8
29	177	1.01	23425	17.4637	4.4	0.70416	6.2	0.089228	4.3	0.706	-1.6	551	23	541.3	25.8	500.7	95.9
30	153	0.79	85703	17.1079	4	0.76137	7.1	0.094512	5.8	0.82	-1	582.2	32.2	574.8	31	545.8	88.3
31	516	0.93	36404	17.1234	3.1	0.68603	4.7	0.085236	3.5	0.75	0.51	527.3	17.8	530.4	19.3	543.8	67.6
33	109	0.75	31031	16.7973	6.3	0.75026	7.9	0.091442	4.8	0.602	0.74	564.1	25.7	568.4	34.4	585.7	136.9
35	353	1.52	41464	17.4222	4	0.69952	4.8	0.088429	2.7	0.557	-1.4	546.2	14.1	538.5	20.2	505.9	88.4
36	962	0.06	47342	18.4788	3.2	0.43383	4.1	0.058169	2.6	0.638	0.35	364.5	9.3	365.9	12.7	374.8	71.3
37	566	0.01	108527	18.1879	3.9	0.51565	6.3	0.068051	4.9	0.785	-0.42	424.4	20.2	422.2	21.6	410.5	86.8
38	91	0.65	22771	17.51	6.5	0.68093	7.9	0.086513	4.6	0.574	-1.4	534.9	23.4	527.3	32.7	494.8	143.4
39	704	0.08	38877	17.6114	2.7	0.57814	3.9	0.073879	2.9	0.735	0.71	459.5	12.8	463.3	14.6	482.1	59
40	145	0.85	39106	17.0627	4.1	0.72045	6.3	0.089196	4.8	0.762	0.024	550.8	25.3	550.9	26.8	551.6	89
41	412	1.54	99985	17.2974	3	0.69957	4.9	0.087803	3.9	0.791	-0.61	542.5	20.2	538.5	20.5	521.6	65.8
43	375	1.37	106457	17.0079	2.7	0.7204	3.9	0.088903	2.8	0.726	0.31	549	14.9	550.9	16.6	558.6	58.5
45	137	0.83	15577	17.0882	7.6	0.56381	9.1	0.069908	5	0.55	4	435.6	21.2	454	33.5	548.3	166.8
46	195	1.20	88539	17.5361	4.3	0.69423	5.8	0.088335	3.8	0.665	-1.8	545.7	20.1	535.3	24	491.5	95
49	170	0.72	57851	17.3756	6.9	0.72171	8.4	0.090991	4.9	0.578	-1.7	561.4	26.1	551.7	35.8	511.8	150.7
50	95	0.95	97635	16.8727	6.7	0.75477	11.3	0.092404	9.2	0.809	0.18	569.7	50.1	571	49.6	576	144.9
51	798	0.06	268948	17.4544	2.4	0.60077	3.8	0.076086	2.9	0.773	0.92	472.7	13.4	477.7	14.4	501.8	52.9
53	161	0.56	12130	17.2139	4.5	0.68991	5.8	0.086173	3.7	0.641	-0.019	532.9	19.2	532.7	24.2	532.3	98.3
55	632	0.02	174434	18.1537	3.4	0.49136	4.6	0.064723	3.1	0.682	0.36	404.3	12.3	405.8	15.4	414.7	75.2
56	427	0.09	49376	18.0132	5.6	0.49549	6.5	0.064762	3.3	0.513	0.97	404.5	13.1	408.6	21.8	432	124.3
58	231	1.09	57695	17.5929	3.8	0.67915	8.5	0.086696	7.6	0.894	-1.5	536	39.2	526.3	35	484.4	84.5
59	416	0.14	115542	17.8793	3.3	0.53677	4.7	0.069636	3.3	0.708	0.48	434	13.8	436.3	16.5	448.6	73
60	527	0.01	104454	18.2736	3.5	0.51504	4.5	0.068291	2.8	0.63	-0.88	425.8	11.7	421.8	15.6	399.9	78.5
61	62	0.56	8119	16.3018	8.2	0.73809	10.7	0.087306	6.9	0.644	3.9	539.6	35.6	561.3	46.1	650.3	175.6
62	164	0.56	42950	12.5943	3.2	2.21153	6.2	0.202098	5.3	0.855	-0.16	1186.6	57.2	1184.7	43.1	1181.3	63.2
65	197	0.54	143755	17.2844	4.5	0.73157	5.6	0.09175	3.4	0.611	-1.4	565.9	18.7	557.5	24.2	523.3	98
67	131	0.79	21675	17.1966	5.3	0.71829	7.4	0.089627	5.2	0.697	-0.6	553.3	27.4	549.7	31.5	534.5	116.5
68	222	0.70	59864	18.3488	5.7	0.53023	6.9	0.070594	3.9	0.567	-1.7	439.7	16.7	432	24.4	390.7	128.4
69	106	0.83	13249	16.9452	5.1	0.72247	7.8	0.088831	5.9	0.758	0.56	548.6	30.9	552.1	33.1	566.6	110.2
71	62	0.70	27743	17.2701	6.6	0.72845	9.7	0.091282	7	0.73	-1.2	563.1	38	555.6	41.3	525.1	144.5
72	243	0.61	19024	17.2405	4	0.69646	6.8	0.087125	5.5	0.809	-0.28	538.5	28.2	536.7	28.1	528.9	87
76	463	0.02	53431	18.4424	3.4	0.49123	4.4	0.065734	2.7	0.622	-1.1	410.4	10.8	405.7	14.6	379.3	77
77	204	0.79	38691	9.2368	2.6	4.60559	5	0.308673	4.3	0.855	1.1	1734.2	65.5	1750.3	42.1	1769.6	47.8
78	674	1.06	149260	17.0776	2.5	0.69922	4.2	0.086644	3.4	0.805	0.41	535.7	17.3	538.3	17.5	549.7	54.2
80	79	0.65	19464	16.7207	6.5	0.71585	8.6	0.086851	5.6	0.648	2	536.9	28.7	548.2	36.4	595.6	141.7
82	249	0.75	25878	17.8493	5.1	0.6075	6.4	0.078679	3.9	0.612	-1.2	488.2	18.4	482	24.6	452.3	112.7
84	124	0.72	25994	17.0306	4.5	0.7183	7.3	0.088763	5.8	0.79	0.22	548.2	30.4	549.7	31.1	555.7	97.9
87	98	0.67	14438	17.2392	7	0.72038	8.5	0.09011	4.9	0.57	-0.92	556.2	25.9	550.9	36.2	529.1	153.1
88	166	1.22	323030	17.2689	4.4	0.69918	7	0.087608	5.4	0.77	-0.48	541.4	27.8	538.3	29.1	525.3	97.3
89	80	0.70	119907	16.9037	5.9	0.74878	7.9	0.091839	5.2	0.661	0.18	566.4	28.2	567.5	34.2	572	128.3
94	114	0.90	36985	16.9669	4.6	0.69217	5.8	0.085214	3.4	0.592	1.3	527.2	17.2	534.1	23.9	563.8	101
96	564	0.34	83903	19.2016	3.4	0.35006	4.5	0.048772	3	0.659	-0.62	307	9	304.8	12	287.8	78
97	359	1.15	113079	17.6949	4.3	0.58958	5.3	0.075697	3	0.565	0.042	470.4	13.5	470.6	19.8	471.6	96
98	438	1.33	39968	17.0193	2.9	0.70079	4.2	0.086542	3	0.708	0.17	535.1	15.2	539.3	17.4	557.1	64.1
99	2444	3.03	120314	17.0222	1.7	0.71259	2.4	0.088013	1.8	0.727	0.4	543.8	9.2	546.3	10.2	556.7	36.2
100	208	0.86	20352	17.0298	4.7	0.67837	6.6	0.083824	4.5	0.693	1.2	518.9	22.7	525.8	26.9	555.8	103.3
103	859	0.01	47320	17.7413	2.5	0.50367	3.6	0.064837	2.5	0.705	2.1	405	9.8	414.2	12.1	465.8	55.9
104	220	1.14	49659	17.234	5	0.71137	6.3	0.088956	3.9	0.616	-0.66	549.4	20.4	545.6	26.5	529.7	108.5
108	156	0.76	35370	17.0379	4.1	0.70317	6.8	0.08693	5.5	0.799	0.52	537.4	28.2	540.7	28.7	554.8	89.8
110	271	1.03	22832	16.8774	4.2	0.73871	5.6	0.090463	3.7	0.661	0.57	558.3	20	561.7	24.4	575.3	92.1
111	1636	1.75	283204	17.214	1.9	0.70921	3.3	0.088584	2.6	0.806	-0.44	547.2	13.8	544.3	13.7	532.3	42.3
112	545	0.02	38835	17.1527	2.9	0.6034	4.4	0.075098	3.3	0.76	2.4	466.8	15	479.4	16.8	540.1	62.4
113	433	1.39	674094	17.2477	2.9	0.68085	4.8	0.085208	3.9	0.807	0.027	527.1	19.7	527.3	19.8	528	62.5
114	271	0.22	11198	17.7051	5.3	0.55865	6.5	0.071768	3.9	0.59	0.81	446.8	16.6	450.6	23.7	470.4	116.6
115	172	0.74	49973	17.0956	4.7	0.71969	6.2	0.089274	4.1	0.65	-0.12	551.2	21.4	550.5	26.5	547.4	103.8
117	333	0.24	46800	17.4861	3.1	0.6651	5.4	0.084386	4.5	0.826	-0.7	522.3	22.4	517.7	22	497.8	67.4
118	76	0.57	14875	17.3294	6.9	0.69316	10.7	0.087159	8.2	0.766	-0.65	538.7	42.5	534.7	44.6	517.6	151.6
120	158	1.03	349816	8.1067	2.2	5.92785	6.1	0.348688	5.6	0.932	2.9	1928.3	94.1	1965.3	52.6	2004.5	39.1
125	343	0.98	62418	17.2344	4	0.68799	4.7	0.086034	2.5	0.522	-0.077	532	12.6	531.6	19.6	529.7	88.6
126	442	0.40	26364	18.263	3.8	0.50712	5.3	0.067201	3.7	0.697	-0.59	419.3	15	416.5	18	401.2	84.8
128	183	1.00	60543	16.9811	4.8	0.69906	6.8	0.086134	4.8	0.705	0.96	532.6	24.6	538.2	28.6	562	105.5
129	410	1.27	184701	17.3145	3.5	0.69086	5	0.086795	3.6	0.718	-0.54	536.6	18.4	533.3	20.7	519.5	76.1
130	526	0.29	1538038	12.6014	2.4	2.16913	4	0.198334	3.1	0.791	0.37	1166.4	33.4	1171.2	27.5	1180.2	47.9
132	225	0.58	22636	10.4311	1.6	3.57845	5.2</										

Table 2. LA-ICP-MS U-Pb zircon data for the Southern Pines 100 K (detrital ages)

Composition			Isotopic Ratios								Ages (Ma)						
ID	U (ppm)	Th/U	²⁰⁶ Pb/ ²⁰⁴ Pb	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (%)	²⁰⁷ Pb/ ²³⁵ U	Error (%)	²⁰⁶ Pb/ ²³⁸ U	Error (%)	error correlation	concordia distance (d _c) ^a	²⁰⁶ Pb/ ²³⁸ U	Error (Ma)	²⁰⁷ Pb/ ²³⁵ U	Error (Ma)	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (Ma)
179	296	0.41	67427	17.422	3.2	0.6839	5.7	0.086454	4.8	0.83	-0.81	534.5	24.5	529.1	23.7	505.9	70.5
181	159	0.88	14677	17.3492	5.6	0.7106	7.3	0.089454	4.8	0.652	-1.2	552.3	25.3	545.1	31	515.1	122.3
182	153	1.04	38913	16.7817	5	0.69682	6.7	0.08485	4.5	0.667	2.1	525	22.7	536.9	28.1	587.7	109.2
183	87	0.91	15895	16.8845	6.7	0.71939	8.5	0.088135	5.2	0.615	1	544.5	27.2	550.3	36	574.4	145.5
184	208	0.05	40190	17.3494	3.5	0.60514	5.2	0.076179	3.8	0.738	1.4	473.3	17.5	480.5	19.9	515.1	77
185	2336	2.08	506430	16.865	1.8	0.71614	2.8	0.087635	2.1	0.763	1.1	541.5	11.1	548.4	11.9	576.9	39.4
186	313	0.54	99011	13.2511	2.1	1.9242	4.4	0.18501	3.9	0.878	-0.47	1094.3	38.9	1089.5	29.4	1080	42.2
187	98	0.65	6796	17.1451	7.9	0.68235	10.1	0.084887	6.4	0.633	0.53	525.2	32.4	528.2	41.8	541	171.7
188	385	1.10	23388	17.3431	3	0.69302	4.8	0.08721	3.8	0.785	-0.68	539	19.7	534.6	20.1	515.9	65.9
189	71	0.67	9871	17.0858	8.2	0.701	11.7	0.086906	8.4	0.715	0.37	537.2	43.1	539.4	48.9	548.6	178.5
190	386	1.59	82379	16.8727	3.5	0.68256	5.1	0.083564	3.6	0.717	1.9	517.4	18.1	528.3	20.9	576	76.8
191	117	0.78	14342	17.5072	6.7	0.69104	8.2	0.087783	4.6	0.562	-1.6	542.4	23.8	533.4	33.8	495.2	148.7
192	430	0.58	61824	17.1618	2.6	0.67517	3.7	0.084075	2.6	0.711	0.6	520.4	13.2	523.8	15.2	538.9	57.3
198	234	1.01	27090	16.9846	3.6	0.71338	4.7	0.087916	3.1	0.652	0.61	543.2	16.1	546.7	20	561.6	78.1
199	141	0.93	12338	16.9916	5.8	0.73497	8	0.090614	5.5	0.687	0.056	559.2	29.4	559.5	34.4	560.7	126.4
200	564	0.02	39642	17.8528	3.6	0.5569	5.1	0.072141	3.6	0.709	0.094	449	15.8	449.5	18.6	451.9	80.4
202	342	0.77	131369	17.5807	3.4	0.6713	4.9	0.085633	3.5	0.719	-1.4	529.7	18	521.5	20	485.9	75.4
203	118	0.92	96690	17.3372	6.1	0.73615	7.6	0.092607	4.5	0.594	-1.8	570.9	24.7	560.2	32.8	516.6	134.6
204	383	1.18	246616	17.2242	4	0.7058	4.5	0.08821	2.2	0.486	-0.49	544.9	11.6	542.3	19.1	531	87.1
205	245	0.94	85251	17.3154	4.3	0.71059	5.5	0.089278	3.4	0.618	-1.1	551.3	17.8	545.1	23.1	519.4	94.4
207	174	0.83	17348	17.2391	5.2	0.73625	6.5	0.092094	4	0.611	-1.3	567.9	21.7	560.2	28.2	529.1	113.5
209	139	0.65	43050	16.9813	4.6	0.70902	7.2	0.087362	5.4	0.762	0.69	539.9	28.2	544.2	30.1	562	100.9
211	220	1.08	51917	17.4449	4.2	0.69672	6.7	0.08819	5.2	0.78	-1.2	544.8	27.1	536.8	27.8	503	91.7
212	331	1.39	157706	17.8137	4.6	0.54183	5.8	0.070035	3.4	0.592	0.69	436.4	14.4	439.6	20.6	456.8	103.1
213	320	0.96	134588	17.2091	3.1	0.69521	4.6	0.08681	3.4	0.736	-0.12	536.6	17.6	535.9	19.4	532.9	69
215	122	0.94	30971	17.3091	6	0.68268	8.2	0.085741	5.6	0.68	-0.33	530.3	28.3	528.4	33.7	520.2	131.5
216	263	1.01	24331	17.3109	3.6	0.69124	5	0.086825	3.5	0.693	-0.54	536.7	17.9	533.5	20.8	519.9	79.2
220	308	1.22	544161	17.1138	4.1	0.69513	7.4	0.086319	6.2	0.834	0.32	533.7	31.8	535.9	31	545.1	89.6
221	180	0.99	19535	17.2122	4.1	0.70401	5.8	0.087924	4.1	0.71	-0.34	543.2	21.6	541.2	24.5	532.5	90
223	202	0.05	113187	17.9075	5.1	0.55098	5.9	0.071592	2.9	0.491	-0.017	445.7	12.4	445.6	21.2	445.1	113.6
224	405	0.79	730776	13.7418	2.7	1.61211	3.5	0.160743	2.2	0.64	1.5	960.9	19.9	974.9	21.9	1006.6	54.3
226	212	0.57	30524	10.7082	2.1	3.29411	4.4	0.255947	3.9	0.884	0.87	1469.1	51.2	1479.7	34.4	1494.9	39.1
228	147	0.96	9427	17.2228	5.1	0.71204	7.1	0.088982	5	0.704	-0.59	549.5	26.4	546	30.1	531.2	110.9
230	150	1.18	7730	17.1081	4.8	0.69097	7	0.085774	5	0.719	0.5	530.5	25.5	533.4	28.9	545.7	105.6
231	83	0.81	16332	17.3279	5.1	0.73835	8.4	0.092833	6.6	0.791	-1.6	572.3	36.3	561.4	36.1	517.8	112.5
232	464	0.02	139084	17.9518	3	0.52604	4.3	0.06852	3	0.705	0.41	427.2	12.5	429.2	14.9	439.6	67.4
233	236	0.80	23475	17.551	4.1	0.70287	5.6	0.08951	3.8	0.68	-2	552.6	20.1	540.5	23.4	489.6	90.3
234	139	0.70	24899	17.1164	5.1	0.70174	8.5	0.087153	6.9	0.806	0.18	538.7	35.6	539.8	35.8	544.7	110.4
236	317	1.47	24495	17.3745	3.6	0.68287	5.2	0.086089	3.7	0.718	-0.65	532.4	18.9	528.5	21.3	511.9	78.9
237	91	0.67	54736	16.8632	4.1	0.72072	6.6	0.088186	5.2	0.786	0.97	544.8	27.2	551.1	28.2	577.2	89.3
240	124	0.67	10622	17.2337	3.9	0.70023	5.6	0.087561	4	0.714	-0.36	541.1	20.7	538.9	23.3	529.8	85.6
242	492	0.04	73115	17.8964	3.6	0.53678	4.9	0.069704	3.3	0.675	0.41	434.4	13.9	436.3	17.4	446.5	80.6
243	85	0.66	82222	17.1718	6.6	0.70452	9	0.087782	6.2	0.687	-0.15	542.4	32.3	541.5	37.9	537.6	143.7
244	109	0.67	21306	17.5582	6.7	0.68566	8.4	0.087354	5.1	0.606	-1.7	539.9	26.3	530.2	34.6	488.7	146.9
245	92	0.74	9130	17.3228	6.9	0.70418	10.1	0.08851	7.3	0.728	-0.88	546.7	38.5	541.3	42.3	518.4	151.7
246	787	0.01	63485	17.8891	2.8	0.5509	4.9	0.071509	4	0.825	0.063	445.2	17.3	445.6	17.6	447.4	61.1
247	128	0.65	7897	17.3279	5.9	0.68582	7.9	0.086228	5.3	0.663	-0.5	533.2	26.9	530.3	32.8	517.8	130.4
248	191	0.88	27406	17.4895	4.2	0.70771	6.5	0.089811	5	0.765	-1.7	554.4	26.5	543.4	27.4	497.4	92.6
249	72	0.75	146373	16.7215	6.5	0.74407	9.5	0.090278	6.8	0.724	1.2	557.2	36.6	564.8	41	595.5	141.4
250	108	0.55	17987	17.0972	7.3	0.72233	10.1	0.08961	6.9	0.687	-0.2	553.2	36.8	552	43	547.2	160.3
253	99	0.66	11668	17.4411	5.8	0.69951	9.2	0.088524	7.1	0.773	-1.3	546.8	37.2	538.5	38.4	503.5	128.3
257	223	0.94	31558	16.8712	3.6	0.70662	5.5	0.086502	4.1	0.75	1.3	534.8	21.2	542.7	23.2	576.2	79.2
258	1041	0.01	48509	17.9331	2.5	0.52807	4	0.068713	3.1	0.785	0.43	428.4	13	430.5	14.1	441.9	55.2
261	606	0.25	320488	16.2859	2.8	0.81351	4.4	0.096132	3.3	0.76	1.9	591.7	18.8	604.4	19.9	652.4	60.9
264	225	0.88	34731	17.2968	5.2	0.70965	6.5	0.089065	4	0.609	-0.94	550	21	544.5	27.6	521.7	114
266	99	0.85	154946	17.1589	8.2	0.70497	10.1	0.087772	6	0.59	-0.099	542.3	31.1	541.8	42.5	539.3	179
267	117	0.63	42433	17.3635	5.6	0.70079	7.2	0.088291	4.5	0.628	-1.1	545.4	23.6	539.3	30	513.3	122.7
268	126	1.02	12035	17.5749	5.3	0.67881	7.2	0.086564	4.8	0.672	-1.6	535.2	24.8	526.1	29.5	486.6	117.6
269	842	1.82	409259	17.2822	2.1	0.72527	3.3	0.090948	2.6	0.78	-1.1	561.1	13.8	553.8	14.1	523.6	45.3
271	445	0.08	84507	16.8984	4.3	0.65937	7.5	0.080848	6.1	0.813	2.2	501.2	29.3	514.2	30.1	572.7	94.5
273	68	0.69	61148	16.5996	7.9	0.7448	10.2	0.089708	6.5	0.635	1.9	553.8	34.4	565.2	44.2	611.3	170.2
275	940	0.16	45823	17.9187	2.8	0.54218	4.1	0.070492	3	0.735	0.15	439.1	12.7	439.9	14.5	443.7	61.4
276	338	0.40	107601	14.8495	3.8	1.17642	5.7	0.126756	4.3	0.747	2.3	769.3	31	789.6	31.4	847.4	79.2
277	415	1.02	44807	17.2708	2.8	0.7044	4.5	0.088272	3.5	0.786	-0.6	545.3	18.5	541.4	18.9	525	61
278	147	0.35	49097	15.7134	3.2	1.04855	5.9	0.119551	5	0.842	0.025	728	34.4	728.2	30.9	728.8	67.8
281	100	0.76	59382	16.9789	5.7	0.73777	7.5	0.090892	4.8	0.646	0.055	560.8	26	561.1	32.3	562.3	124.7
283	569	0.48	206350	15.3901	2.9	1.15143	7	0.12858	6.4	0.912	-0.26	779.8	46.9	777.9	38.1	772.6	60.4
284	323	0.85	24741	17.4435	4.7	0.71474	6.2	0.090464	4.1	0.66	-1.8	558.3	21.9	547.6	26.3	503.2	102.6
287	93	0.67	1														

Table 2. LA-ICP-MS U-Pb zircon data for the Southern Pines 100 K (detrital ages)

Composition			Isotopic Ratios									Ages (Ma)							
ID	U (ppm)	Th/U	²⁰⁶ Pb/ ²⁰⁴ Pb			²⁰⁶ Pb/ ²⁰⁷ Pb			error			concordia		²⁰⁶ Pb/ ²³⁸ U		²⁰⁷ Pb/ ²³⁵ U		²⁰⁶ Pb/ ²⁰⁷ Pb	
			Value	Error (%)	2σ	Value	Error (%)	2σ	Value	Error (%)	2σ	distance (d _c) ^a	Age	Error (Ma)	Value	Error (Ma)	Age	Error (Ma)	
26	171	0.93	9950	16.933	5.9	0.69661	6.8	0.085589	3.3	0.493	1.4	529.4	17	536.8	28.2	568.2	128.2		
29	124	0.85	89727	16.8363	4.2	0.75551	5.1	0.092296	3	0.588	0.39	569.1	16.5	571.4	22.5	580.7	90.4		
31	75	0.85	9227	17.1787	5.2	0.7034	6.9	0.087677	4.5	0.657	-0.17	541.8	23.5	540.8	28.9	536.7	113.7		
32	133	0.72	24839	17.437	2.3	0.68121	3.8	0.086188	3.1	0.796	-0.84	533	15.6	527.5	15.8	504	51.1		
33	90	0.43	27339	17.2735	4.8	0.69596	5.9	0.087229	3.4	0.582	-0.49	539.1	17.7	536.4	24.5	524.7	105		
34	153	0.64	24071	17.3082	3.3	0.71179	5.6	0.089392	4.6	0.81	-0.9	551.9	24.2	545.8	23.8	520.3	72.5		
36	190	0.58	13921	17.4346	3.1	0.71619	4.5	0.090601	3.2	0.718	-1.7	559.1	17.2	548.4	19	504.3	68.7		
37	203	0.77	95157	17.2333	2.8	0.71623	4	0.08956	2.9	0.713	-0.72	552.9	15.2	548.4	17.1	529.8	61.9		
42	300	0.55	78339	16.9923	2.8	0.73242	3.6	0.090303	2.3	0.626	0.11	557.3	12.1	558	15.5	560.6	61.4		
45	88	0.78	28829	17.3807	4	0.70346	5.4	0.088716	3.6	0.673	-1.2	547.9	19.1	540.9	22.7	511.1	88		
49	607	0.96	29178	16.6539	3.5	0.73209	4.4	0.088466	2.8	0.625	2	546.5	14.5	557.8	19	604.2	74.9		
50	271	0.79	17589	17.0472	3.1	0.70609	4.2	0.087339	2.9	0.687	0.45	539.8	15	542.4	17.7	553.6	66.9		
52	183	0.78	13607	16.385	3.3	0.7512	4.1	0.089309	2.4	0.586	3.1	551.4	12.6	568.9	17.7	639.4	71		
54	144	0.64	250903	17.0615	2.9	0.7104	4.4	0.087945	3.3	0.749	0.25	543.4	17.1	545	18.5	551.7	63.5		
57	149	0.67	75987	17.1006	3.5	0.71819	4.7	0.089113	3.1	0.666	-0.11	550.3	16.5	549.6	20	546.7	76.8		
62	60	0.60	19071	17.1123	5.1	0.72214	6.6	0.089666	4.1	0.629	-0.29	553.6	22	551.9	28.1	545.2	112.1		
63	186	0.53	11050	17.3571	4.1	0.69329	5.1	0.087314	3	0.591	-0.86	539.6	15.6	534.8	21.2	514.1	90.2		
64	1047	0.70	137012	16.8588	1.5	0.70832	2.3	0.086647	1.8	0.763	1.3	535.7	9.2	543.7	9.8	577.7	32.8		
65	516	0.53	487445	17.0102	2.8	0.70852	3.7	0.08745	2.5	0.668	0.58	540.4	12.9	543.9	15.7	558.3	60.4		
67	284	0.75	12237	17.3902	3.4	0.6958	3.9	0.087798	2	0.511	-1.1	542.5	10.5	536.3	16.4	509.9	74.3		
68	288	0.82	53502	17.2321	2.9	0.7023	4.1	0.087811	2.9	0.716	-0.4	542.6	15.3	540.2	17.2	530	62.6		
Sample 550946																			
2	112	0.40	87034	17.4781	4.1	0.70459	5.4	0.089356	3.4	0.636	-1.8	551.7	18	541.5	22.5	498.8	91.2		
4	249	0.78	21778	17.0961	3.8	0.66054	4.6	0.081938	2.6	0.562	1.4	507.7	12.8	514.9	18.8	547.3	84		
5	207	0.88	9089	16.806	8.8	0.70853	10.1	0.0864	4.9	0.487	1.8	534.2	25.3	543.9	42.6	584.6	191.7		
7	218	0.85	18156	17.6355	2.9	0.6849	4.5	0.087641	3.4	0.753	-1.9	541.6	17.5	529.7	18.4	479	65		
9	284	0.69	33336	17.2032	3.3	0.69364	4.6	0.086584	3.2	0.696	-0.054	535.3	16.3	535	18.9	533.6	71.7		
13	198	0.63	34178	17.3189	4.2	0.69158	5	0.086907	2.8	0.556	-0.63	537.2	14.5	533.7	21	518.9	92.2		
15	206	0.54	22811	17.4336	4.2	0.68034	5.5	0.086061	3.6	0.659	-0.92	532.2	18.6	527	22.7	504.4	91.5		
16	167	0.60	141050	17.3995	3.3	0.6481	9.9	0.081822	9.4	0.944	0.075	507	45.8	507.3	39.7	508.7	72.1		
17	122	0.72	6961	17.4323	7	0.68919	8.2	0.087174	4.2	0.513	-1.2	538.8	21.8	532.3	34	504.6	155.1		
18	288	0.71	28197	17.5028	2.9	0.68875	3.9	0.08747	2.6	0.66	-1.4	540.6	13.3	532	16.1	495.7	64.5		
19	123	0.73	47393	17.3625	4.2	0.71273	6	0.089791	4.3	0.722	-1.3	554.3	23	546.4	25.4	513.4	91.4		
22	540	1.01	135300	16.9069	2.1	0.68601	6	0.084157	5.6	0.938	1.9	520.9	28.2	530.4	24.8	571.5	45.4		
24	178	0.52	381241	16.9112	2.8	0.6966	4.8	0.085478	3.9	0.806	1.3	528.7	19.5	536.8	19.9	571	61.6		
25	91	0.69	60559	17.3569	4	0.68243	5.9	0.085946	4.4	0.737	-0.54	531.5	22.3	528.2	24.4	514.1	87.9		
26	254	0.66	35449	17.5617	2.6	0.68517	3.5	0.087308	2.3	0.663	-1.7	539.6	11.9	529.9	14.3	488.3	57.3		
27	197	0.87	10508	16.7929	5.5	0.68171	7.1	0.083065	4.4	0.629	2.5	514.4	21.9	527.8	29	586.2	118.9		
29	340	0.78	16242	16.5515	4.6	0.70463	5.7	0.084623	3.4	0.601	3.3	523.7	17.2	541.5	24	617.6	98.5		
33	208	0.50	44463	17.347	2.7	0.66726	3.8	0.083987	2.7	0.696	-0.14	519.9	13.2	519	15.5	515.4	60.1		
36	165	0.79	34859	17.3885	2.6	0.69723	3.8	0.08797	2.8	0.744	-1	543.5	14.8	537.1	16	510.1	56.1		
40	315	1.19	77630	17.373	2.1	0.6737	3.8	0.084925	3.1	0.823	-0.39	525.5	15.6	523	15.3	512.1	46.9		
41	172	0.57	71534	17.423	3.2	0.71183	5.6	0.089989	4.6	0.822	-1.4	555.5	24.6	545.8	23.7	505.8	70.3		
42	129	0.58	10309	17.6266	5.1	0.67558	6.5	0.086405	3.9	0.606	-1.8	534.2	20.1	524.1	26.5	480.1	113.7		
45	206	0.65	63036	17.2174	2.8	0.69828	4.7	0.087236	3.8	0.809	-0.22	539.2	19.9	537.8	19.8	531.8	61.1		
46	129	0.57	20084	17.5054	4	0.70803	5.3	0.089932	3.4	0.643	-2	555.1	18	543.6	22.2	495.4	89.1		
50	284	0.67	12047	17.4727	2.3	0.67442	3.5	0.085503	2.7	0.765	-0.88	528.9	13.7	523.4	14.4	499.5	50		
51	127	0.53	16529	17.4198	3.5	0.69792	5.9	0.088216	4.7	0.8	-1.1	545	24.7	537.6	24.6	506.2	77.9		
52	139	0.51	13323	17.5147</															

Table 2. LA-ICP-MS U-Pb zircon data for the Southern Pines 100 K (detrital ages)

Composition			Isotopic Ratios						Ages (Ma)								
ID	U (ppm)	Th/U	²⁰⁶ Pb/ ²⁰⁴ Pb	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (%)	²⁰⁷ Pb/ ²³⁵ U	Error (%)	²⁰⁶ Pb/ ²³⁸ U	Error (%)	error correlation	concordia distance (d _c) ^a	²⁰⁶ Pb/ ²³⁸ U	Error (Ma)	²⁰⁷ Pb/ ²³⁵ U	Error (Ma)	²⁰⁶ Pb/ ²⁰⁷ Pb	Error (Ma)
146	267	0.83	86327	17.222	2.5	0.68861	3.6	0.08605	2.6	0.714	-0.029	532.1	13.3	532	15.1	531.3	55.9
148	232	0.60	18938	17.3585	3.7	0.68794	4.6	0.086648	2.7	0.579	-0.75	535.7	13.6	531.6	18.9	513.9	82
149	161	0.74	21873	17.3622	3.3	0.67142	4.3	0.084585	2.7	0.625	-0.34	523.4	13.4	521.6	17.5	513.4	73.4
151	385	0.70	45832	17.1566	2.3	0.70337	3.7	0.08756	2.9	0.78	-0.042	541.1	15.2	540.8	15.7	539.6	51.3
154	291	1.01	28750	17.4039	2.3	0.65846	4	0.083151	3.3	0.824	-0.19	514.9	16.3	513.7	16.1	508.2	49.6
155	130	0.48	40283	17.5438	4.9	0.68343	6.3	0.086998	4	0.636	-1.6	537.8	20.7	528.8	26	490.5	107.3
159	162	0.87	23310	17.3435	3.3	0.68339	5	0.086	3.7	0.752	-0.5	531.8	19	528.8	20.4	515.8	71.8
160	149	0.56	16500	17.601	3.9	0.67221	5.5	0.08585	3.9	0.712	-1.5	530.9	19.9	522.1	22.4	483.4	85.2
161	659	1.10	134731	17.1709	2.3	0.72261	4.2	0.090031	3.5	0.841	-0.51	555.7	18.8	552.2	17.8	537.7	49.5
162	186	0.65	127808	17.4364	3.1	0.69636	4.2	0.088101	2.8	0.672	-1.3	544.3	14.8	536.6	17.6	504.1	68.7
163	228	0.79	59846	17.1391	2.6	0.71287	4	0.088653	3	0.758	-0.17	547.6	15.8	546.5	16.8	541.8	56.6
167	191	0.59	37467	17.4563	2.6	0.68847	5.8	0.087203	5.2	0.891	-1.1	539	26.9	531.9	24.2	501.6	58.3
168	119	0.51	43962	17.5647	3.7	0.67071	5.3	0.08548	3.8	0.721	-1.3	528.8	19.3	521.1	21.5	487.9	80.7
169	118	0.66	5942	16.5319	6.8	0.71411	8.3	0.08566	4.8	0.576	3.2	529.8	24.4	547.2	35.2	620.1	146.7
171	202	0.94	15573	16.7191	3.8	0.74855	5	0.090809	3.2	0.648	1.2	560.3	17.3	567.4	21.6	595.8	82
172	204	0.64	63503	17.3782	3.1	0.68179	4.3	0.085971	3	0.699	-0.65	531.7	15.5	527.9	17.9	511.4	68.4
173	170	0.56	51605	17.4976	3.4	0.68477	5.2	0.08694	4	0.765	-1.2	537.4	20.5	529.7	21.5	496.3	73.9
174	151	0.53	249909	17.2968	3.4	0.70105	4.8	0.087985	3.5	0.715	-0.68	543.6	18	539.4	20.2	521.7	74.2
178	147	0.55	16043	17.5617	3.9	0.66863	5.2	0.085201	3.4	0.651	-1.3	527.1	17.1	519.9	21.1	488.3	86.7
179	120	0.53	11891	17.6027	4.8	0.68006	5.8	0.08686	3.2	0.552	-1.8	536.9	16.4	526.8	23.7	483.2	106
181	167	0.52	61590	17.2937	3.1	0.70198	4.2	0.088086	2.9	0.675	-0.71	544.2	14.9	540	17.8	522.1	68.7
182	123	0.53	62342	17.2017	3.8	0.70027	5.2	0.087403	3.6	0.691	-0.2	540.2	18.7	539	21.8	533.8	82.6
183	147	0.56	64812	17.1781	3.5	0.70154	5.5	0.087442	4.2	0.765	-0.11	540.4	21.8	539.7	23	536.8	77.5
184	194	0.66	284438	17.5654	3	0.69235	4.3	0.088242	3.1	0.719	-1.8	545.1	16	534.2	17.7	487.8	65.2
185	162	0.56	63235	17.4254	2	0.69119	3.3	0.087392	2.6	0.796	-1	540.1	13.5	533.5	13.6	505.5	43.5
186	134	0.68	11956	17.6937	3.8	0.67544	6.4	0.086716	5.1	0.803	-1.9	536.1	26.5	524	26.2	471.7	84.6
188	200	0.59	513440	17.4089	2.4	0.6872	4.5	0.086805	3.8	0.846	-0.83	536.6	19.4	531.1	18.4	507.6	52.2
189	191	0.55	24380	17.4944	3.1	0.71101	4.6	0.090255	3.5	0.748	-1.8	557	18.5	545.3	19.5	496.8	67.5
191	179	0.49	17652	17.5136	3.6	0.6854	5.7	0.087099	4.5	0.782	-1.3	538.4	23.2	530	23.7	494.3	78.9
192	176	0.68	9113	17.426	4.9	0.70578	5.6	0.08924	2.9	0.51	-1.6	551	15.2	542.2	23.7	505.4	106.8
195	79	0.66	14011	17.4478	5.7	0.65203	7.4	0.082548	4.7	0.64	-0.3	511.3	23.2	509.7	29.6	502.7	125
196	220	0.64	39559	17.4289	2.5	0.71239	4.2	0.090092	3.4	0.805	-1.5	556.1	18	546.2	17.7	505	54.7
199	170	0.53	171301	17.018	3.7	0.69464	4.9	0.085775	3.1	0.645	0.9	530.5	16	535.6	20.3	557.3	81.3
201	173	0.57	14424	17.5352	4.4	0.68943	6.3	0.08772	4.5	0.719	-1.6	542	23.4	532.5	26	491.6	96.1
203	149	0.54	76730	17.443	3.2	0.6839	4.6	0.086558	3.3	0.714	-1	535.1	16.8	529.1	18.9	503.2	70.5
205	47	0.67	15170	16.765	5.6	0.68955	9.3	0.083881	7.4	0.8	2.2	519.2	37.1	532.5	38.6	589.9	121.3
208	155	0.57	29573	17.3292	3.8	0.69346	5	0.087196	3.2	0.646	-0.71	538.9	16.8	534.9	20.9	517.6	84.3
210	195	0.77	74753	16.8582	2.2	0.72035	4.4	0.088115	3.8	0.862	0.95	544.4	19.6	550.9	18.5	577.8	48
214	308	1.01	29461	16.8792	2.8	0.66728	4	0.081725	2.9	0.719	2.2	506.4	14	519.1	16.2	575.1	60.2
218	193	0.94	54466	17.3454	2.7	0.70496	4.8	0.088724	4	0.831	-0.92	548	21.1	541.7	20.3	515.6	59.1
219	159	0.71	46533	17.1047	4	0.70494	5.9	0.087491	4.3	0.731	0.17	540.7	22.2	541.7	24.6	546.2	87.2
223	208	0.54	30170	17.1164	3.1	0.6955	4.7	0.086379	3.6	0.759	0.32	534.1	18.2	536.1	19.5	544.7	66.7
224	144	0.89	45754	17.5995	3.4	0.68041	5.1	0.086889	3.8	0.743	-1.6	537.1	19.3	527	20.8	483.6	74.7
225	194	0.54	123513	16.7807	3.5	0.71027	5.3	0.086482	3.9	0.741	1.7	534.7	20.1	544.9	22.3	587.8	77
228	170	0.53	139136	17.3834	2.9	0.6688	6	0.084358	5.2	0.875	-0.33	522.1	26.2	520	24.3	510.8	63.5
229	122	0.76	7756	17.5936	6.1	0.68816	7.4	0.087849	4.2	0.566	-2	542.8	21.8	531.7	30.7	484.3	134.8
231	185	0.62	47581	17.2199	2.4	0.70448	3.8	0.088023	2.9	0.777	-0.37	543.8	15.3	541.5	15.8	531.5	51.9
233	145	0.47	54967	17.6624	2.9	0.66823	4.6	0.085638	3.5	0.772	-1.6	529.7	17.9	519.6	18.5	475.7	64
234	221	0.76	41870	17.1654	2	0.68753	3.7	0.085633	3.1	0.839	0.25	529.7	15.9	531.3	15.4	538.4	44.5
235	317	0.93	158781	17.2693	2.5	0.69689	3.9	0.087324	3	0.768	-0.43	539.7	15.4	536.9	16.1	525.2	54.3
236	142	0.50	27665	17.7123	3.6	0.66106	5.3	0.084959	3.9	0.733	-1.7	525.7	19.6	515.3	21.4	469.4	79.7
237	141	0.79	10123	17.4236	5.5	0.6666	6.7	0.084275	3.8	0.572	-0.55	521.6	19.1	518.6	27.1	505.7	120.5
240	153	0.80	48063	17.245	4.4	0.69023	6.2	0.086368	4.3	0.696	-0.19	534	22.1	532.9	25.7	528.3	97.3
243	306	1.02	83593	17.2066	2.8	0.73097	3.9	0.091262	2.8	0.699	-0.92	563	14.9	557.1	16.9	533.2	61.7
244	146	0.49	13610	17.5114	4.5	0.67881	6.4	0.08625	4.5	0.703	-1.2	533.3	23	526.1	26.2	494.6	100.2
248	151	0.81	118557	17.223	3.8	0.67695	5.6	0.084598	4	0.721	0.25	523.5	20.1	524.9	22.8	531.1	84.3
249	127	0.62	13904	17.4267	4.6	0.68394	6.1	0.086483	3.9	0.65	-0.98	534.7	20.2	529.2	25	505.3	101.6
251	132	0.72	208845	16.8385	2.7	0.70173	3.9	0.085737	2.8	0.728	1.6	530.3	14.4	539.8	16.2	580.4	57.7
252	140	0.51	43929	17.1448	3.1	0.68942	4.7	0.085765	3.6	0.756	0.32	530.4	18.1	532.5	19.5	541.1	67.2
255	93	0.34	6449	17.5196	7.1	0.64726	8.3	0.08228	4.2	0.509	-0.56	509.7	20.6	506.8	32.9	493.6	156.8
259	177	0.50	41057	17.2858	3.9	0.67742	6.3	0.084965	5	0.791	-0.082	525.7	25.2	525.2	25.9	523.1	84.8
262	206	0.69	16086	17.1094	4.2	0.70614	5.5	0.087664	3.5	0.645	0.13	541.7	18.3	542.5	22.9	545.6	91
266	133	0.76	25796	17.4683	3.9	0.67977	5.3	0.08616	3.6	0.675	-1.1	532.8	18.4	526.6	21.9	500.1	86.6
268	233	0.56	11201	17.537	2.7	0.68806	3.8	0.087554	2.6	0.702	-1.6	541.1	13.7	531.6	15.5	491.4	58.9
271	159	0.80	19443	17.5681	3.3	0.67927	4.9	0.086588	3.5	0.726	-1.5	535.3	18.1	526.3	19.9	487.5	73.6
272	117	0.56	44060	17.1303	4.1	0.67615	6.1	0.084043	4.6	0.749	0.72	520.2	23	524.4	25.1	542.9	88.7
273	101	0.67	204398	17.325	3.8	0.73605	5.8	0.092528	4.5	0.767	-1.5	570.5	24.5	560.1	25.2	518.1	82.4
275	197	0.54	50473	17.3304	3.1	0.68304	4.3	0.085891	3	0.686	-0.44	531.2	15.1	528.6	17.8	517.5	68.8
277	111	0.80	30244	17.5812	4.2	0.68814</											