

**FISHERY MANAGEMENT PLAN UPDATE
BLUEFISH
AUGUST 2025**

STATUS OF THE FISHERY MANAGEMENT PLAN

Fishery Management Plan History

FMP Documentation:	1990	
	Amendment 1	2000
	Framework 1	2001
	Amendment 2	2007
	Amendment 3	2011
	Addendum I	2012
	Amendment 4	2013
	Amendment 5	2015
	Amendment 6	2017
	Framework 2	2017
	Framework 3	2018
	Framework 4	2020
	Framework 5	2020
	Amendment 7	2021
	Framework 6	2023
Comprehensive Review:	2023	

The Fishery Management Plan (FMP) for bluefish was developed through a joint management effort between the interstate Atlantic States Marine Fisheries Commission (ASMFC) and the federal Mid-Atlantic Fishery Management Council (MAFMC). Amendment 1 initiated a 10-year rebuilding schedule to eliminate overfishing and allow for stock rebuilding which was achieved in 2009. Amendment 1 also established commercial and recreational quota allocations, state-specific commercial allocations, and allowed for the transfer of unused recreational quota to the commercial fishery. Framework 1 established annual harvest allocations specifically for biological monitoring programs. Amendments 2 and 5 were implemented to establish a strategy for monitoring bluefish bycatch. Amendment 3 added a formalizing process to incorporate scientific and management uncertainty when establishing catch limits. Addendum I established a coast-wide biological monitoring program to improve the quantity and quality of information available for use in bluefish stock assessments. Amendment 4 modified the accountability measures for the recreational bluefish fishery. Amendment 6 addressed considerations for examining potential influence of the removal of forage fish species by increasing directed fishing and advocated for future ecosystem-based management approaches. Framework 2 required for-hire vessels with federal permits for species managed by MAFMC to submit electronic vessel trip reports to the National Oceanic and Atmospheric Administration. Framework 3 established a process to specify constant multi-year acceptable biological catches. Framework 4 established a requirement for commercial vessels with federal permits for any species managed by the Mid-Atlantic and New England Councils to submit vessel trip reports electronically within 48 hours after entering port at the conclusion of a trip. Framework 5 modified the Council's acceptable biological catch control rule and risk policy. The revised risk policy is intended to reduce the probability of overfishing as stock size falls below the target biomass while allowing for increased risk and greater economic benefit under higher stock biomass conditions. This action also removed the typical/atypical species distinction currently included in the risk policy. Amendment 7, the Bluefish Allocation and Rebuilding Amendment, revised the goals and objectives of the fishery management plan, reallocated quota between the commercial and recreational fisheries, reallocated commercial quota among the states,

implemented a rebuilding plan, revised the sector quota transfer process, and revised how management uncertainty is applied during the specifications process. Amendment 7 took effect on January 1, 2022. Framework 6 established a new process for setting recreational bag, size, and season limits for summer flounder, scup, black sea bass, and bluefish. This action also modified recreational accountability measures for these species. Framework 6 took effect on March 9, 2023. The bluefish FMP, associated amendment documents, and framework information can be found at [MSFMC.org](https://www.msfmc.org).

To ensure compliance with interstate requirements, North Carolina also manages bluefish under the North Carolina Fishery Management Plan for Interjurisdictional Fisheries (IJ FMP). The goal of the IJ FMP is to adopt fishery management plans consistent with N.C. law and approved by the MAFMC, South Atlantic Fishery Management Council, or the ASMFC by reference and implement corresponding fishery regulations in North Carolina to provide compliance or compatibility with approved fishery management plans and amendments, now and in the future. The goal of these plans, established under the Magnuson-Stevens Fishery Conservation and Management Act (federal council plans) and the Atlantic Coastal Fisheries Cooperative Management Act (ASMFC plans), are like the goals of the Fisheries Reform Act of 1997 to “ensure long-term viability” of these fisheries (NCDMF 2022a).

Management Unit

The FMP defines the management unit of bluefish as a single stock occurring in U.S. waters of the western Atlantic Ocean. All member Atlantic states participate in the ASMFC bluefish FMP process except for Pennsylvania and the District of Columbia.

Goal and Objectives

Amendment 7 revised the goals and objectives of the bluefish FMP to the following:

- Goal 1: Conserve the bluefish resource through stakeholder engagement to maintain sustainable recreational fishing and commercial harvest.
 - Objective 1.1: Achieve and maintain a sustainable spawning stock biomass and rate of fishing mortality.
 - Objective 1.2: Promote practices that reduce release mortality within the recreational and commercial fishery.
 - Objective 1.3: Maintain effective coordination between the National Marine Fisheries Service, Council, Commission, and member states by promoting compliance and to support the development and implementation of management measures.
 - Objective 1.4: Promote compliance and effective enforcement of regulations.
 - Objective 1.5: Promote science, monitoring, and data collection that support and enhance effective ecosystem-based management of the bluefish resource.
- Goal 2: Provide fair and equitable access to the fishery across all user groups throughout the management unit.
 - Objective 2.1: Ensure the implementation of management measures provides fair and equitable access to the resource across all user groups within the management unit.
 - Objective 2.2: Consider the economic and social needs and priorities of all groups that access the bluefish resource in the development of new management measures.
 - Objective 2.3: Maintain effective coordination with stakeholder groups to ensure optimization of economic and social benefits.

DESCRIPTION OF THE STOCK

Biological Profile

Bluefish (*Pomatomus saltatrix*) are a migratory, open water (pelagic) species found throughout the Atlantic Ocean. Bluefish migrate seasonally, moving north as water temperatures rise during spring and summer and south during the fall and winter to areas along the South Atlantic Bight (Shepherd et al. 2006). During the summer, bluefish mostly concentrate in waters from Maine to Cape Hatteras (Klein-MacPhee 2002). During the winter, they are found in offshore waters between North Carolina and Florida (Goodbred and Graves 1996). Within North Carolina's estuarine waters, bluefish are most common from March through October. Bluefish generally school with similarly sized fish (Austin et al. 1999). Bluefish are fast growers (Wilk 1977) and opportunistic predators. Over 70 different marine species have been documented in bluefish stomach contents including Atlantic menhaden, butterfish, silversides, spotted seatrout, Atlantic croaker, spot, shrimp, lobster, squid, crabs, worms, and clams (Buckel et al. 1999; Scharf et al. 2004). The maximum documented age for bluefish is 14 years (Robillard et al. 2009). Bluefish can exceed 39 inches and 31 pounds (NCDMF 2022b). Bluefish usually reach sexual maturity by age two around a length of 13 inches (Robillard et al. 2008). They spawn offshore from Massachusetts through Florida. Bluefish born each year typically fall into two distinct size classes, suggesting that there are two distinct spawning events, with one group spawning during the spring and a second spawning during the summer (Lassiter 1962). However, more recent research suggests that bluefish spawning is a single, continuous event that occurs as they migrate northward during the spring and summer, but that bluefish spawned in the middle of this time period do not have high survivability, resulting in two distinct size groups (Smith et al. 1994; Robillard et al. 2008).

Stock Status

The 2023 management track stock assessment determined that bluefish are not overfished and are not experiencing overfishing.

Stock Assessment

Results from the 2023 management track assessment indicate that the Atlantic bluefish stock was not overfished and not experiencing overfishing in 2022. SSB in 2022 was estimated at 0.152, or 64% of the overfishing threshold of 0.239. Although fishing mortality was below the threshold in 2022, fishing mortality exceeded the updated threshold every year from 1985 to 2017, except 2008. Recruitment has increased annually since 2019 but has remained below the time series average over the past 12 years, except in 2022. The next management track assessment for bluefish is scheduled for 2025.




Figure 1. Bluefish spawning stock biomass and recruitment at age 0 by calendar year. The horizontal dashed red line is the SSB Threshold = 100,865 mt.

DESCRIPTION OF THE FISHERY

Current Regulations

In North Carolina, the private recreational (all persons not fishing on a for-hire vessel) bag limit is three bluefish per person per day and the recreational for-hire (all persons fishing on a for-hire vessel) is five bluefish per person per day. These regulations have been in effect since 2020. Commercial fishery landings are monitored and if necessary, trip limits are implemented to prevent exceeding the annual quota. The commercial fishery was opened on January 1, 2024, with no possession limit. Commercial possession limits were decreased three times during 2024: 800-pound limit on February 22, 400-pound limit on May 21, 50-pound limit and then increased to a 300-pound limit on September 6.

Commercial Fishery

In North Carolina, bluefish have been harvested commercially using a variety of gears including estuarine long haul, ocean trawl, pound net, ocean beach seine, ocean gill net, and estuarine gill net. Capture methods have shifted primarily to gill nets over the last few decades. Gill nets, especially estuarine gill nets, have been the primary mode of harvest. Estuarine and ocean gill nets combined represent the largest commercial landings of bluefish, accounting for ~96% of the harvest in 2024 (Figure 2).




Figure 2. Commercial harvest of bluefish in North Carolina during 2024 by gear type.

The commercial quota allocated to North Carolina for 2024 was 776,452 pounds. Additionally, North Carolina received a total of 394,544 pounds of quota transfers from Massachusetts, New York, New Jersey, Maryland, and Virginia totaling 1,170,996 pounds. North Carolina's 2024 commercial bluefish landings totaled 1,193,181 pounds at a dockside value of \$745,588. Bluefish commercial landings have fluctuated annually since 1985 (Figure 3); however, landings in 2024 decreased slightly from 2023 (Table 1).




Figure 3. North Carolina commercial landings of bluefish, 1985–2024.

Table 1. Bluefish recreational harvest and number released (Marine Recreational Information Program) and commercial harvest (North Carolina Trip Ticket Program) in North Carolina, 2015–2024.

Year	Recreational			Commercial	
	Number Landed	Number Released	Weight Landed (lb)	Weight Landed (lb)	Total Weight Landed (lb)
2015	4,123,461	6,356,252	3,754,577	804,847	4,559,424
2016	4,489,223	6,802,960	3,356,049	1,148,643	4,504,692
2017	3,173,218	8,255,510	3,634,502	1,544,053	5,178,555
2018	3,304,587	7,912,210	2,630,685	910,262	3,540,947
2019	2,752,589	7,162,431	3,011,480	1,108,205	4,119,685
2020	2,108,296	6,557,751	2,124,224	1,112,966	3,237,190
2021	982,389	3,539,333	1,031,760	1,051,019	2,082,779
2022	1,533,911	9,336,045	1,645,410	872,042	2,517,452
2023	1,261,404	4,775,374	1,492,689	1,658,869	3,151,558
2024	1,574,579	6,390,035	2,512,747	1,193,181	3,705,928
Mean	2,535,818	6,724,129	2,510,379	1,140,409	3,650,788

There is a wide distribution of bluefish harvested in the commercial fishery, however the majority of commercially caught bluefish are between 12 and 18 inches (Figure 4). In the last decade, bluefish longer than 20 inches are harvested less often than compared to the rest of the time series (Figure 5B).




Figure 4. Commercial and recreational length frequency distribution from bluefish harvested in North Carolina, 2024.




Figure 5. Recreational (A) and commercial (B) length frequency (fork length, inches) of bluefish harvested from 1985–2024. Bubbles represent fish at length and the bubble size is proportional to the number of fish at that length.

Recreational Fishery

Bluefish are caught recreationally from shore, pier, and boat and can be targeted with lures as well as live and dead bait. Discards are a large part of the Bluefish fishery as they are not highly sought after for eating.

In 2024, approximately 80% of landed fish were released (Table 1). Recreational landings of bluefish vary annually but have declined in the last decade (Figure 6).




Figure 6. North Carolina recreational landings of bluefish, 1985–2024.

In 2024, the size distribution of fish taken in the recreational fishery was similar to the distribution of fish harvested in the commercial fishery (Figure 4). However, the percentage of bluefish harvested at each size was less consistent across the distribution (Figure 5A). For bluefish exceeding 15 pounds or 34 inches, the NCDMF offers award citations. The number of citations awarded was highest in 1991 ($n=187$), with fewer citations awarded in the last 20 years (Figure 7). Approximately 61% of the citations awarded since 2017 have been for released fish.




Figure 7. North Carolina recreational award citations for bluefish, 2005–2024.

MONITORING PROGRAM DATA

Fishery-Dependent Monitoring

Commercial bluefish landings from a broad range of gears are sampled through the Division's fish house sampling programs. Information collected includes location, gear type and gear-specifics, soak time, and water depth. Commercial catches are also subsampled to collect biological information on bluefish including fork length (FL) and aggregate weight (kg) by market grade. Trip ticket information (total weight of catch) is also recorded and reported to DMF by licensed dealers. A total of 2,672 bluefish were measured from commercial landings in 2024 (Table 2). Mean fork length was 17 inches and ranged from 9 to 32 inches. Mean size and size ranges have varied minimally over the last few decades. Since 1985, the mean size of bluefish landed is 13 inches fork length with a mean minimum fork length of 6 inches and a mean maximum fork length of 34 inches.

The number and size of fish harvested as well as number of fish released recreationally is characterized through NOAA Fisheries' Marine Recreational Information Program (MRIP). In 2024, approximately 2.5 million pounds of bluefish were recreationally harvested (Table 1). The mean length of fish harvested and measured by MRIP in the recreational fishery in 2024 was 14 inches and ranged from 8 to 32 inches fork length (Table 2). Since 1985, the annual length distribution of harvest in both the commercial and recreational fisheries has varied little with most fish harvested ranging from 7 to 16 inches fork length (Figure 5). Larger bluefish (>20 inches) have been less common in recent years in both the commercial and recreational fisheries. See [NOAA](#) for more information on the collection of recreational fishing data.

Table 2. Summary of fork length (inches) data sampled from all sources of length data (harvest and bait) from the bluefish commercial fishery and the bluefish recreational fishery in North Carolina, 2004–2024.

Year	Commercial				Recreational			
	Mean Length	Minimum Length	Maximum Length	Total Number Measured	Mean Length	Minimum Length	Maximum Length	Total Number Measured
2004	19	6	33	9,608	13	6	40	1,149
2005	19	5	33	9,766	12	6	35	1,056
2006	18	5	33	10,255	12	6	36	1,028
2007	15	6	33	8,856	12	6	37	1,048
2008	16	5	33	8,035	12	5	35	894
2009	18	6	34	7,471	13	7	34	778
2010	17	6	35	6,721	12	6	38	1,323
2011	16	6	33	5,768	12	6	34	1,784
2012	14	5	34	7,030	12	7	35	1,190
2013	14	6	33	6,928	11	7	29	563
2014	15	8	34	6,459	12	7	29	660
2015	14	7	31	6,100	12	7	18	577
2016	14	3	33	7,616	11	8	23	732
2017	16	7	35	5,580	12	6	35	657
2018	15	7	34	3,778	11	6	30	846
2019	15	8	33	4,812	13	8	32	910
2020	16	7	35	3,396	12	8	32	713
2021	16	8	34	4,203	12	6	26	299
2022	14	4	31	3,945	12	8	29	433
2023	15	7	29	4,701	13	7	29	413
2024	17	9	32	2,672	14	8	32	269

Fishery-Independent Monitoring

The Division's Pamlico Sound Independent Gill Net Survey was initiated in May of 2001 and has been sampled continuously throughout 2019. Sampling in this program was suspended in February 2020 due to COVID-19 restrictions - but resumed continuous sampling July 2021. This survey provides fishery-independent indices of relative abundance along with associated length and age data. The relative abundance index, defined as the number of bluefish per set, provides essential data for input into the coastwide bluefish stock assessment. The relative abundance index in 2024 was 13.8, which is more than double the time-series average of 6.1 (Figure 8). The 2023 and 2024 abundance indices are the highest in the time series, with the lowest being 2.8 in 2015. It should be noted that the index in 2021 is calculated from samples collected from Jul–Dec while the index for all other years was calculated for Feb–Dec.




Figure 8. Relative abundance index of bluefish, from the North Carolina Pamlico Sound Independent Gill Net Survey, 2001–2024. Shading represents the standard error about the annual relative abundance index estimates.

North Carolina is one of the states subject to compliance of the biological monitoring program implemented under Addendum I to Amendment 1. To comply with these monitoring requirements, DMF must collect at least 100 ageing structures from bluefish each year with at least 50 fish collected from January–June and 50 fish from July–December. Most bluefish age samples are collected from the Pamlico Sound Independent Gill Net Survey. Other age sample sources include commercial and recreational fisheries. In 2024, 998 age samples were collected (Table 3). The maximum age in 2024 was 8 years of age. The maximum age observed in the time-series is 12 years. Bluefish length increases with age, although the size at a given age is variable (Figure 9).

Table 3. Summary of bluefish age samples collected in North Carolina from both dependent and independent sources, 2009–2024.

Year	Modal Age	Min. Age	Max. Age	Number of Samples
2009	3	0	10	488
2010	3	0	8	527
2011	3	0	9	551
2012	1	0	9	818
2013	0	0	9	742
2014	1	0	9	803
2015	1	0	10	622
2016	1	0	11	678
2017	2	0	10	630
2018	1	0	10	669
2019	1	0	8	853
2020	2	0	12	244
2021	1	0	5	793
2022	1	0	8	1,210
2023	1	0	6	1,170
2024	2	0	8	998




Figure 9. Bluefish length at age based on all age samples collected in North Carolina, 1985–2024. Blue circles represent the mean size at a given age while the grey squares represent the minimum and maximum observed size for each age.

RESEARCH NEEDS

- Evaluate magnitude and length frequency of discards from the commercial and recreational fisheries, especially recreational discard lengths in the mid-Atlantic and southern regions

- Develop additional adult bluefish indices of abundance (e.g., broad spatial and temporal scale longline survey or gill-net survey) to better characterize dynamics of older bluefish not well sampled by fisheries-independent surveys
- Explore age- and time-varying natural mortality from, for example, predator-prey relationships; quantify effects of age- and time-varying mortality on the assessment model
- Investigate potential spatial distribution shifts of the Atlantic stock
- Initiate coastal surf zone seine study to provide more complete indices of juvenile abundance.
- Expand age structure of Southeast Area Monitoring and Assessment Program index.
- Investigate species associations with recreational angler trips targeting bluefish (on a regional and seasonal basis) to potentially modify the MRIP index used in the assessment model.
- Continue to evaluate the spatial, temporal, and sector-specific trends in bluefish growth and quantify their effects in the assessment model.
- Continue to examine alternative models that take advantage of length-based assessment frameworks.
- Evaluate the source of bimodal length frequency in the catch (e.g., migration, differential growth rates).
- Modify thermal niche model to incorporate water temperature data more appropriate for bluefish in a timelier manner [e.g., sea surface temperature data & temperature data that cover the full range of bluefish habitat (South Atlantic Bight and estuaries)].
- Investigate potential spatial distribution shifts of the Atlantic stock.

MANAGEMENT

Bluefish in North Carolina are jointly managed by ASMFC and MAFMC under Amendment 2 of the FMP. Amendment 2 uses annual catch limits (ACLs) for both the recreational and commercial sectors. The recreational quota is a coast-wide quota while the commercial quota is further divided into state-specific quotas. Amendment 2 allows quota transfers between states and between sectors. Additionally, daily limits are used to manage recreational harvest and trip limits can be implemented for commercial fishermen if needed to prevent exceeding North Carolina's commercial quota.

LITERATURE CITED

- Austin, H.M., D. Scoles, and A.J. Abell. 1999. Morphometric separation of annual cohorts within mid-Atlantic bluefish, *Pomatomus saltatrix*, using discriminant function analysis. *Fishery Bulletin*. 97(3):411-420
- Buckel, J.A., M.J. Fogarty, and D.O. Conover. 1999. Foraging habits of bluefish, *Pomatomus saltatrix*, on the U.S. east coast continental shelf. *Fishery Bulletin*. 97:758-775
- Goodbred, C.O., and J.E. Graves. 1996. Genetic relationships among geographically isolated populations of bluefish (*Pomatomus saltatrix*). *Marine and Freshwater Research* 47:347-355
- Klein-MacPhee, G. 2002. Bluefish: family Pomatomidae. In Bigelow and Schroeder's fishes of the Gulf of Maine (B.B. Collette, and G. Klein-MacPhee, eds.), p. 400-406. Smithsonian Institution Press, Washington, D.C.
- Lassiter, R.R. 1962. Life history aspects of the bluefish, *Pomatomus saltatrix*, larvae and juveniles off the east coast of the United States. *Fishery Bulletin*. 77:213-227
- NCDMF (North Carolina Division of Marine Fisheries). 2022a. North Carolina Fishery Management Plan for Interjurisdictional Fisheries, 2022 Information Update. North Carolina Division of Marine Fisheries, Morehead City, North Carolina. 19 pp.

- NCDMF. 2022b. North Carolina State Saltwater Records. <https://deq.nc.gov/about/divisions/marine-fisheries/public-information-and-education/coastal-fishing-information/nc-saltwater-fishing-tournament/north-carolina-state-saltwater-records>
- NMFS, Northeast Fisheries Science Center. 2021. Atlantic Bluefish Operational Assessment for 2021. US Department of Commerce, Northeast Fisheries Science Center. Woods Hole, Massachusetts.
- Robillard, E., C.S. Reiss, and C.M. Jones. 2008. Reproductive biology of bluefish (*Pomatomus saltatrix*) along the east coast of the United States. Fisheries Research. 90:198-208
- Robillard, E., C.S. Reiss, and C.M. Jones. 2009. Age-validation and growth of bluefish (*Pomatomus saltatrix*) along the east coast of the United States. Fisheries Research. 95(1): 65-75
- Scharf, F.S., J.P. Manderson, M.C. Fabrizio, J.P. Pessutti, J.E. Rosendale, R.J. Chant, and A.J. Bejda. 2004. Seasonal and interannual patterns of distribution and diet of bluefish within a Middle Atlantic Bight estuary in relation to abiotic and biotic factors. Estuaries, 27(3): 428-436
- Shepherd, G.R., J. Moser, D. Deuel, and P. Carlson. 2006. The migration patterns of bluefish (*Pomatomus saltatrix*) along the Atlantic coast determined from tag recoveries. Fishery Bulletin. 104:559-570
- Smith, W., P. Berrien, and T. Potthoff. 1994. Spawning patterns of bluefish, *Pomatomus saltatrix*, in the northeast continental shelf ecosystem. Bull. Mar. Sci. 54(1): 8-16.
- Wilk, S.J. 1977. Biological and fisheries data on bluefish, *Pomatomus saltatrix* (Linnaeus). NOAA, NMFS, NEFC, Sandy Hook Lab. Technical Series Report. No. 11.